

EXACT SOLUTION OF THE SIX-VERTEX MODEL WITH DOMAIN WALL BOUNDARY CONDITIONS. ANTIFERROELECTRIC PHASE

PAVEL BLEHER AND KARL LIECHTY

ABSTRACT. We obtain the large n asymptotics of the partition function Z_n of the six-vertex model with domain wall boundary conditions in the antiferroelectric phase region, with the weights $a = \sinh(\gamma - t)$, $b = \sinh(\gamma + t)$, $c = \sinh(2\gamma)$, $|t| < \gamma$. We prove the conjecture of Zinn-Justin, that as $n \rightarrow \infty$, $Z_n = C\vartheta_4(n\omega)F^{n^2}[1 + O(n^{-1})]$, where ω and F are given by explicit expressions in γ and t , and $\vartheta_4(z)$ is the Jacobi theta function. The proof is based on the Riemann-Hilbert approach to the large n asymptotic expansion of the underlying discrete orthogonal polynomials and on the Deift-Zhou nonlinear steepest descent method.

1. INTRODUCTION AND FORMULATION OF THE MAIN RESULT

1.1. Definition of the model. The six-vertex model, or the model of two-dimensional ice, is stated on a square $n \times n$ lattice with arrows on edges. The arrows obey the rule that at every vertex there are two arrows pointing in and two arrows pointing out. Such a rule is sometimes called the *ice-rule*. There are only six possible configurations of arrows at each vertex, hence the name of the model, see Fig. 1.




FIGURE 1. The six arrow configurations allowed at a vertex.

Date: December 16, 2009.

The first author is supported in part by the National Science Foundation (NSF) Grant DMS-0652005.

Both authors would like to thank the Centre de Recherches Mathématiques at l'Université de Montréal, where much of this work was performed, for its hospitality during their visit in the Fall of 2008.

We will consider the *domain wall boundary conditions* (DWBC), in which the arrows on the upper and lower boundaries point into the square, and the ones on the left and right boundaries point out. One possible configuration with DWBC on the 4×4 lattice is shown on Fig. 2.




FIGURE 2. An example of 4×4 configuration with DWBC.

For each possible vertex state we assign a weight w_i , $i = 1, \dots, 6$, and define, as usual, the partition function as a sum over all possible arrow configurations of the product of the vertex weights,

$$Z_n = \sum_{\text{arrow configurations } \sigma} w(\sigma), \quad w(\sigma) = \prod_{x \in V_n} w_{t(x; \sigma)} = \prod_{i=1}^6 w_i^{N_i(\sigma)}, \quad (1.1)$$

where V_n is the $n \times n$ set of vertices, $t(x; \sigma) \in \{1, \dots, 6\}$ is the vertex-type of configuration σ at vertex x according to Fig. 1, and $N_i(\sigma)$ is the number of vertices of type i in the configuration σ . The sum is taken over all possible configurations obeying the given boundary condition. The Gibbs measure is defined then as

$$\mu_n(\sigma) = \frac{w(\sigma)}{Z_n}. \quad (1.2)$$

Our main goal is to obtain the large n asymptotics of the partition function Z_n .

The six-vertex model has six parameters: the weights w_i . By using some conservation laws it can be reduced to only two parameters (see, e.g., [14], [7], and [1]). Namely, we have that

$$Z_n(w_1, w_2, w_3, w_4, w_5, w_6) = C(n) Z_n(a, a, b, b, c, c), \quad (1.3)$$

and

$$\mu_n(\sigma; w_1, w_2, w_3, w_4, w_5, w_6) = \mu_n(\sigma; a, a, b, b, c, c), \quad (1.4)$$

where

$$a = \sqrt{w_1 w_2}, \quad b = \sqrt{w_3 w_4}, \quad c = \sqrt{w_5 w_6}, \quad (1.5)$$

and

$$C(n) = \left(\frac{w_5}{w_6} \right)^{\frac{n}{2}}. \quad (1.6)$$

Furthermore,

$$Z_n(a, a, b, b, c, c) = c^{n^2} Z_n \left(\frac{a}{c}, \frac{a}{c}, \frac{b}{c}, \frac{b}{c}, 1, 1 \right) \quad (1.7)$$

and

$$\mu_n(\sigma; a, a, b, b, c, c) = \mu_n \left(\sigma; \frac{a}{c}, \frac{a}{c}, \frac{b}{c}, \frac{b}{c}, 1, 1 \right), \quad (1.8)$$

so that a general weight reduces to the two parameters, $\frac{a}{c}, \frac{b}{c}$.

1.2. Exact solution of the six-vertex model for a finite n . Introduce the parameter

$$\Delta = \frac{a^2 + b^2 - c^2}{2ab}. \quad (1.9)$$

The phase diagram of the six-vertex model consists of the three phase regions: the ferroelectric phase region, $\Delta > 1$; the antiferroelectric phase region, $\Delta < -1$; and the disordered phase region, $-1 < \Delta < 1$. Observe that $|a - b| > c$ in the ferroelectric phase region and $c > a + b$ in the antiferroelectric phase region, while in the disordered phase region a, b, c satisfy the triangle inequalities.




FIGURE 3. The phase diagram of the model, where **F**, **AF** and **D** mark ferroelectric, antiferroelectric, and disordered phases, respectively. The circular arc corresponds to the so-called free fermion line, when $\Delta = 0$, and the three dots correspond to 1-, 2-, and 3-enumeration of alternating sign matrices.

In the three phase regions we parameterize the weights in the standard way: for the ferroelectric phase,

$$a = \sinh(t - \gamma), \quad b = \sinh(t + \gamma), \quad c = \sinh(2|\gamma|), \quad 0 < |\gamma| < t; \quad (1.10)$$

for the antiferroelectric phase,

$$a = \sinh(\gamma - t), \quad b = \sinh(\gamma + t), \quad c = \sinh(2\gamma), \quad |t| < \gamma; \quad (1.11)$$

and for the disordered phase

$$a = \sin(\gamma - t), \quad b = \sin(\gamma + t), \quad c = \sin(2\gamma), \quad |t| < \gamma. \quad (1.12)$$

The phase diagram of the six-vertex model is shown on Fig. 3. The phase diagram and the Bethe Ansatz solution of the six-vertex model for periodic and antiperiodic boundary conditions are thoroughly discussed in the works of Lieb [21]-[24], Lieb, Wu [25], Sutherland [28], Baxter [2], Batchelor, Baxter, O'Rourke, Yung [3]. See also the work of Wu, Lin [29], in which the Pfaffian solution for the six-vertex model with periodic boundary conditions is obtained on the free fermion line, $\Delta = 0$. Brascamp, Kunz and Wu [6] prove the equality of the free energy with periodic and free boundary conditions under various conditions on a, b, c , and also they prove the existence of the spontaneous staggered polarization for sufficiently small values of the parameters a/c and b/c .

As concerns the six-vertex model with DWBC, it was noticed by Kuperberg [20], that on the diagonal,

$$\frac{a}{c} = \frac{b}{c} = x, \quad (1.13)$$

the six-vertex model with DWBC is equivalent to the s -enumeration of alternating sign matrices (ASM), in which the weight of each such matrix is equal to s^{N_-} , where N_- is the number of (-1) 's in the matrix and $s = \frac{1}{x^2}$. The exact solution for a finite n is known for 1-, 2-, and 3-enumerations of ASMs, see the works by Kuperberg [20] and Colomo-Pronko [11] for a solution based on the Izergin-Korepin formula. A fascinating story of the discovery of the ASM formula is presented in the book [10] of Bressoud. On the free fermion line, $\gamma = \frac{\pi}{4}$, the partition function of the six-vertex model with DWBC has a very simple form: $Z_n = 1$. For a nice short proof of this formula see the work [11] of Colomo-Pronko.

In this paper we will discuss the *antiferroelectric phase region*, and we will use parameterization (1.11). The parameter Δ in the antiferroelectric phase region reduces to

$$\Delta = -\cosh(2\gamma). \quad (1.14)$$

The six-vertex model with DWBC was introduced by Korepin in [17], who derived important recursion relations for the partition function of the model. These recursion relations have been solved by Izergin [15], and this lead to a beautiful determinantal formula for the partition function with DWBC. A detailed proof of this formula, usually called the Izergin-Korepin formula, and its generalizations are given in the paper of Izergin, Coker, and Korepin [16]. When the weights are parameterized according to (1.11), the Izergin-Korepin formula is

$$Z_n = \frac{[\sinh(\gamma - t) \sinh(\gamma + t)]^{n^2} \tau_n}{\left(\prod_{j=0}^{n-1} j!\right)^2}, \quad (1.15)$$

where τ_n is the Hankel determinant,

$$\tau_n = \det \left(\frac{d^{j+k-2} \phi}{dt^{j+k-2}} \right)_{1 \leq j, k \leq n}, \quad (1.16)$$

and

$$\phi(t) = \frac{\sinh(2\gamma)}{\sinh(\gamma - t) \sinh(\gamma + t)}. \quad (1.17)$$

An elegant derivation of the Izergin-Korepin formula from the Yang-Baxter equations is given in the papers of Korepin, Zinn-Justin [18] and Kuperberg [20] (see also the book of Bressoud [10]).

One of the applications of the determinantal formula is that it implies that the function τ_n solves the Toda equation

$$\tau_n \tau_n'' - \tau_n'^2 = \tau_{n+1} \tau_{n-1}, \quad n \geq 1, \quad (') = \frac{\partial}{\partial t}, \quad (1.18)$$

cf. the work of Sogo, [27]. The Toda equation was used by Korepin and Zinn-Justin [18] to derive the free energy of the six-vertex model with DWBC, assuming some ansatz on the behavior of subdominant terms in the large n asymptotics of the free energy.

Another application of the Izergin-Korepin formula is that τ_n can be expressed in terms of a partition function of a random matrix model and also in terms of related orthogonal polynomials, see the paper [31] of Zinn-Justin. In the antiferroelectric phase the expression in terms of orthogonal polynomials can be obtained as follows. For the evaluation of the Hankel determinant, let us write $\phi(t)$ in the form of the Laplace transform of a discrete measure,

$$\phi(t) = \frac{\sinh(2\gamma)}{\sinh(\gamma - t) \sinh(\gamma + t)} = 2 \sum_{l=-\infty}^{\infty} e^{2tl - 2\gamma|l|}. \quad (1.19)$$

Then

$$\tau_n = \frac{2^{n^2}}{n!} \sum_{l_1, \dots, l_n = -\infty}^{\infty} \Delta(l)^2 \prod_{i=1}^n e^{2tl_i - 2\gamma|l_i|}, \quad (1.20)$$

where

$$\Delta(l) = \prod_{1 \leq i < j \leq n} (l_j - l_i) \quad (1.21)$$

is the Vandermonde determinant, see Appendix A in the end of the paper.

Introduce now discrete monic polynomials $P_j(x) = x^j + \dots$ orthogonal on the set $\mathbb{Z}$ with respect to the weight,

$$w(l) = e^{2tl - 2\gamma|l|}, \quad (1.22)$$

so that

$$\sum_{l=-\infty}^{\infty} P_j(l) P_k(l) w(l) = h_k \delta_{jk}. \quad (1.23)$$

Then it follows from (1.20) that

$$\tau_n = 2^{n^2} \prod_{k=0}^{n-1} h_k, \quad (1.24)$$

see Appendix B in the end of the paper.

1.3. **Rescaling of the weight.** Set

$$\Delta_n = \frac{2\gamma}{n}, \quad x = l\Delta_n, \quad w_n(x) = e^{-n(|x|-\zeta x)}, \quad \zeta = \frac{t}{\gamma}, \quad (1.25)$$

and

$$P_{nk}(x) = \Delta_n^k P_k\left(\frac{x}{\Delta_n}\right). \quad (1.26)$$

Consider also the lattice

$$L_n = \left\{ x = \frac{2\gamma k}{n}, \quad k \in \mathbb{Z} \right\}. \quad (1.27)$$

Then from (1.23) we obtain that the monic polynomials $P_{nk}(x)$ satisfy the orthogonality condition,

$$\sum_{x \in L_n} P_{nj}(x) P_{nk}(x) w_n(x) \Delta_n = h_{nk} \delta_{jk}, \quad h_{nk} = h_k \Delta_n^{2k+1}. \quad (1.28)$$

We can then combine equations (1.15), (1.24), and (1.28) to obtain

$$Z_n = \left(\frac{nab}{\gamma} \right)^{n^2} \prod_{k=0}^{n-1} \frac{h_{nk}}{(k!)^2}, \quad a = \sinh(\gamma - t), \quad b = \sinh(\gamma + t). \quad (1.29)$$

For what follows we will need to extend the weight $w_n(x)$ to the complex plane. We do so by defining $w_n(z)$ on the complex plane as

$$w_n(z) = e^{-nV(z)} \quad (1.30)$$

where

$$V(z) = \begin{cases} z - \zeta z & \text{when } \operatorname{Re} z \geq 0, \\ -z - \zeta z & \text{when } \operatorname{Re} z \leq 0, \end{cases} \quad (1.31)$$

so that $V(z)$, and thus $w_n(z)$, is two-valued on the imaginary axis.

1.4. **Main result: Asymptotics of the partition function.** This work is a continuation of the work [4] of the first author with Vladimir Fokin and [7], [8] of the authors of the present work. In [4] the authors obtain the large n asymptotics of the partition function Z_n in the disordered phase. They prove the conjecture of Paul Zinn-Justin [31] that the large n asymptotics of Z_n in the disordered phase has the following form: For some $\varepsilon > 0$,

$$Z_n = C n^\kappa F^{n^2} [1 + O(n^{-\varepsilon})]. \quad (1.32)$$

Furthermore, they find the exact value of the exponent κ ,

$$\kappa = \frac{1}{12} - \frac{2\gamma^2}{3\pi(\pi - 2\gamma)}. \quad (1.33)$$

The value of F in the disordered phase is given by the formula,

$$F = \frac{\pi ab}{2\gamma \cos \frac{\pi t}{2\gamma}}, \quad a = \sin(\gamma - t), \quad b = \sin(\gamma + t), \quad (1.34)$$

in parameterization (1.12).

In the work [7] we obtain the following large n asymptotic formula for Z_n in the ferroelectric phase region: For any $\varepsilon > 0$,

$$Z_n = CG^n F^{n^2} \left[1 + O\left(e^{-n^{1-\varepsilon}}\right) \right], \quad (1.35)$$

where $C = 1 - e^{-4\gamma}$, $G = e^{\gamma-t}$, and $F = \sinh(t + \gamma)$ in parameterization (1.10).

Finally, in the work [8] we obtain the following large n asymptotic formula for Z_n on the border line between the ferroelectric and disordered phase regions:

$$Z_n(a, a, a+1, a+1, 1, 1) = Cn^\kappa G^{\sqrt{n}} F^{n^2} [1 + O(n^{-1/2})], \quad (1.36)$$

where $C > 0$,

$$\kappa = \frac{1}{4}, \quad G = \exp \left[-\zeta \left(\frac{3}{2} \right) \sqrt{\frac{a}{\pi}} \right], \quad F = a + 1, \quad (1.37)$$

and ζ is the Riemann zeta-function.

In the present paper we obtain the large n asymptotic formula for Z_n in the antiferroelectric phase region. The formulation of the main result of the present paper and the proofs involve the Jacobi theta functions. Let us review their definition and basic properties.

There are four Jacobi theta functions:

$$\begin{aligned} \vartheta_1(z) &= 2 \sum_{n=0}^{\infty} (-1)^n q^{(n+\frac{1}{2})^2} \sin((2n+1)z) \\ \vartheta_2(z) &= 2 \sum_{n=0}^{\infty} q^{(n+\frac{1}{2})^2} \cos((2n+1)z) \\ \vartheta_3(z) &= 1 + 2 \sum_{n=1}^{\infty} q^{n^2} \cos(2nz) \\ \vartheta_4(z) &= 1 + 2 \sum_{n=1}^{\infty} (-1)^n q^{n^2} \cos(2nz) \end{aligned} \quad (1.38)$$

where q is the *elliptic nome*. We will assume that $1 > q > 0$. Fig. 4 shows the graphs of ϑ_1, ϑ_2 (left figure) and ϑ_3, ϑ_4 (right figure) on the interval $[0, \pi]$ for $q = 0.5$. Observe that ϑ_1, ϑ_4 are increasing on $[0, \frac{\pi}{2}]$ while ϑ_2, ϑ_3 are decreasing on this interval.

The Jacobi theta functions satisfy the following periodicity conditions:

$$\begin{aligned} \vartheta_1(z + \pi) &= -\vartheta_1(z), & \vartheta_1(z + \pi\tau) &= -e^{-2iz} q^{-1} \vartheta_1(z), \\ \vartheta_2(z + \pi) &= -\vartheta_2(z), & \vartheta_2(z + \pi\tau) &= e^{-2iz} q^{-1} \vartheta_2(z), \\ \vartheta_3(z + \pi) &= \vartheta_3(z), & \vartheta_3(z + \pi\tau) &= e^{-2iz} q^{-1} \vartheta_3(z), \\ \vartheta_4(z + \pi) &= \vartheta_4(z), & \vartheta_4(z + \pi\tau) &= -e^{-2iz} q^{-1} \vartheta_4(z), \end{aligned} \quad (1.39)$$

where τ is a pure imaginary number related to q by the equation,

$$q = e^{i\pi\tau}. \quad (1.40)$$

The theta functions also satisfy the symmetry conditions:

$$\vartheta_1(-z) = -\vartheta_1(z), \quad \vartheta_2(-z) = \vartheta_2(z), \quad \vartheta_3(-z) = \vartheta_3(z), \quad \vartheta_4(-z) = \vartheta_4(z), \quad (1.41)$$




FIGURE 4. The graphs of ϑ_1, ϑ_2 (left figure) and ϑ_3, ϑ_4 (right figure) on the interval $[0, \pi]$ for $q = 0.5$.

and the equations,

$$\vartheta_1(z) = \vartheta_2\left(z - \frac{\pi}{2}\right), \quad \vartheta_3(z) = \vartheta_4\left(z + \frac{\pi}{2}\right), \quad \vartheta_1(z) = -ie^{iz + \frac{i\pi\tau}{4}}\vartheta_4\left(z + \frac{\pi\tau}{2}\right). \quad (1.42)$$

The only zeroes of the theta functions are

$$\vartheta_1(0) = 0, \quad \vartheta_2\left(\frac{\pi}{2}\right) = 0, \quad \vartheta_3\left(\frac{\pi}{2} + \frac{\pi\tau}{2}\right) = 0, \quad \vartheta_4\left(\frac{\pi\tau}{2}\right) = 0, \quad (1.43)$$

and their shifts by $m\pi + n\pi\tau$; $m, n \in \mathbb{Z}$. There are many non-trivial identities satisfied by the theta functions. A list of those identities used in this paper is given in Appendix E.

In the antiferroelectric phase region we use parameterization (1.11) with two parameters t, γ such that $|t| < \gamma$. In what follows we will also use the following parameters:

$$\zeta = \frac{t}{\gamma} \in (-1, 1), \quad \omega = \frac{\pi(1 + \zeta)}{2} \in (0, \pi). \quad (1.44)$$

The elliptic nome for all Jacobi theta functions in this paper will be equal to

$$q = e^{-\frac{\pi^2}{2\gamma}}. \quad (1.45)$$

Our main result in the present paper is the following asymptotic formula for Z_n :

Theorem 1.1. *As $n \rightarrow \infty$,*

$$Z_n = C\vartheta_4(n\omega)F^{n^2}(1 + O(n^{-1})), \quad (1.46)$$

where $C > 0$ is a constant, and

$$F = \frac{\pi \sinh(\gamma - t) \sinh(\gamma + t) \vartheta_1'(0)}{2\gamma \vartheta_1(\omega)}. \quad (1.47)$$

The asymptotic formula (1.46) proves the conjecture of Zinn-Justin in [31]. The proof of Theorem 1.1 will be based on the Riemann-Hilbert approach to discrete orthogonal polynomials. An important first step in this approach is a construction of the equilibrium measure.

2. EQUILIBRIUM MEASURE

2.1. Heuristic motivation and definition of the equilibrium measure. If we scale the variables in (1.20) as $\mu_i = \frac{2\gamma l_i}{n}$, then we can rewrite formula (1.20) as

$$\tau_n = \frac{2^{n^2}}{n!} \sum_{\mu \in \frac{2\gamma}{n}\mathbb{Z}^n} e^{-n^2 H_n(\nu_\mu)}, \quad (2.1)$$

where

$$d\nu_\mu(x) = \frac{1}{n} \sum_{j=1}^n \delta(x - \mu_j), \quad (2.2)$$

and

$$H(\nu) = \iint_{x \neq y} \log \frac{1}{|x - y|} d\nu(x) d\nu(y) + \int (|x| - \zeta x) d\nu(x). \quad (2.3)$$

where all integrals are over $\mathbb{R}$.

Due to the factor $(-n^2)$ in the exponent of (2.1), we expect the sum, in the large n limit, to be focused in a neighborhood of a global minimum of the functional H . Clearly, we have that ν_μ is a probability measure and

$$\nu_\mu(a, b) \leq \frac{b - a}{2\gamma} \quad \text{for any} \quad -\infty < a < b < \infty, \quad (2.4)$$

because in (2.2), $\mu_j \in \frac{2\gamma}{n}\mathbb{Z}$ and $\mu_j \neq \mu_k$ if $j \neq k$. With these constraints in mind, we define

$$E_0 = \inf_{\nu} H(\nu) \quad (2.5)$$

where the infimum is taken over all probability measures satisfying (2.4). It is possible to prove that there exists a unique minimizer ν_0 , so that

$$E_0 = H(\nu_0), \quad (2.6)$$

see, e.g., the works of Saff and Totik [26], Dragnev and Saff [13] and Kuijlaars [19]. Furthermore, ν_0 has support on a finite number of intervals, and is absolutely continuous with respect to the Lebesgue measure. The minimizer ν_0 is called the *equilibrium measure*.

Denote the density function of the equilibrium measure as $\rho(x)$, and its resolvent as ω , so we have

$$\frac{d\nu_0}{dx} = \rho(x), \quad \omega(z) = \int \frac{\rho(x) dx}{z - x}, \quad (2.7)$$

and

$$\rho(x) = \frac{1}{2\pi i} (\omega(x - i0) - \omega(x + i0)). \quad (2.8)$$

The structure of the equilibrium measure ν_0 is studied in the paper of Zinn-Justin [31], who shows that ν_0 has support on an interval $[\alpha, \beta]$, with a saturated region $[\alpha', \beta']$ in which

$$\rho(x) = \frac{1}{2\gamma}, \quad x \in [\alpha', \beta'], \quad (2.9)$$

and two unsaturated regions, $[\alpha, \alpha']$ and $[\beta', \beta]$, in which

$$0 < \rho(x) < \frac{1}{2\gamma}, \quad x \in (\alpha, \alpha') \cup (\beta', \beta), \quad (2.10)$$

see Fig. 5. We also have that

$$\alpha < \alpha' < 0 < \beta' < \beta, \quad (2.11)$$

so that the origin, which is a singular point of the potential $V(x) = |x| - \zeta x$, lies inside the saturated region $[\alpha', \beta']$.




FIGURE 5. The equilibrium density function $\rho(x)$.

The measure ν_0 is uniquely determined by the Euler-Lagrange variational conditions

$$2 \int \log |x - y| d\nu_0(y) - (|x| - \zeta x) \begin{cases} = l & \text{for } x \in [\alpha, \alpha'] \cup [\beta', \beta], \\ \geq l & \text{for } x \in [\alpha', \beta'], \\ \leq l & \text{for } x \notin [\alpha, \beta], \end{cases} \quad (2.12)$$

where l is the Lagrange multiplier. The Euler-Lagrange variational conditions imply

$$\omega(x - i0) + \omega(x + i0) = -\zeta + \operatorname{sgn}(x) \quad \text{for } x \in [\alpha, \alpha'] \cup [\beta', \beta], \quad (2.13)$$

whereas in the saturated region, we have

$$\rho(x) = \frac{1}{2\pi i} (\omega(x - i0) - \omega(x + i0)) = \frac{1}{2\gamma} \quad \text{for } x \in [\alpha', \beta']. \quad (2.14)$$

Now we will give a detailed description of the equilibrium measure. We begin with explicit formulae for the end-points of the support of the equilibrium measure.

2.2. Explicit formulae for the end-points. They are given in the following proposition.

Proposition 2.1. *The end-points of the support of the equilibrium measure ν_0 are equal to*

$$\begin{aligned} \alpha &= -\pi \frac{\vartheta'_1(\frac{\omega}{2})}{\vartheta_1(\frac{\omega}{2})}, & \alpha' &= -\pi \frac{\vartheta'_4(\frac{\omega}{2})}{\vartheta_4(\frac{\omega}{2})}, \\ \beta' &= -\pi \frac{\vartheta'_3(\frac{\omega}{2})}{\vartheta_3(\frac{\omega}{2})}, & \beta &= -\pi \frac{\vartheta'_2(\frac{\omega}{2})}{\vartheta_2(\frac{\omega}{2})}. \end{aligned} \quad (2.15)$$

The differences between the end-points are equal to

$$\begin{aligned}\alpha' - \alpha &= \pi \vartheta_4^2(0) \frac{\vartheta_2(\frac{\omega}{2}) \vartheta_3(\frac{\omega}{2})}{\vartheta_1(\frac{\omega}{2}) \vartheta_4(\frac{\omega}{2})}, & \beta' - \alpha' &= \pi \vartheta_2^2(0) \frac{\vartheta_1(\frac{\omega}{2}) \vartheta_2(\frac{\omega}{2})}{\vartheta_3(\frac{\omega}{2}) \vartheta_4(\frac{\omega}{2})}, \\ \beta - \beta' &= \pi \vartheta_4(0)^2 \frac{\vartheta_1(\frac{\omega}{2}) \vartheta_4(\frac{\omega}{2})}{\vartheta_2(\frac{\omega}{2}) \vartheta_3(\frac{\omega}{2})}.\end{aligned}\tag{2.16}$$

and

$$\begin{aligned}\beta - \alpha &= \pi \vartheta_2^2(0) \frac{\vartheta_3(\frac{\omega}{2}) \vartheta_4(\frac{\omega}{2})}{\vartheta_1(\frac{\omega}{2}) \vartheta_2(\frac{\omega}{2})}, & \beta - \alpha' &= \pi \vartheta_3(0)^2 \frac{\vartheta_1(\frac{\omega}{2}) \vartheta_3(\frac{\omega}{2})}{\vartheta_2(\frac{\omega}{2}) \vartheta_4(\frac{\omega}{2})}, \\ \beta' - \alpha &= \pi \vartheta_3^2(0) \frac{\vartheta_2(\frac{\omega}{2}) \vartheta_4(\frac{\omega}{2})}{\vartheta_1(\frac{\omega}{2}) \vartheta_3(\frac{\omega}{2})}.\end{aligned}\tag{2.17}$$

Finally, we have the Zinn-Justin formula for the centroid of the end-points,

$$\frac{\alpha + \alpha' + \beta' + \beta}{4} = -\frac{\pi}{2} \frac{\vartheta_2'(\frac{\pi\zeta}{2})}{\vartheta_2(\frac{\pi\zeta}{2})}.\tag{2.18}$$

For a proof of Proposition 2.1 see the next section.

2.3. Equilibrium density function. The equilibrium density function is described in the following proposition.

Proposition 2.2. *The equilibrium density function $\rho(x)$ is given by the formulae,*

$$\rho(x) = \begin{cases} \frac{1}{\pi} \int_{\alpha}^x \frac{dx'}{\sqrt{(x' - \alpha)(\alpha' - x')(\beta' - x')(\beta - x')}}, & \alpha \leq x \leq \alpha', \\ \frac{1}{2\gamma}, & \alpha' \leq x \leq \beta', \\ \frac{1}{\pi} \int_x^{\beta} \frac{dx'}{\sqrt{(x' - \alpha)(x' - \alpha')(x' - \beta')(\beta - x')}}, & \beta' \leq x \leq \beta. \end{cases}\tag{2.19}$$

Also,

$$\int_0^{\beta} \rho(x) dx = \frac{1 + \zeta}{2}.\tag{2.20}$$

The resolvent $\omega(z)$ of the equilibrium measure is given as

$$\omega(z) = \int_z^{\infty} \frac{dz'}{\sqrt{(z' - \alpha)(z' - \alpha')(z' - \beta')(z' - \beta)}},\tag{2.21}$$

where integration takes place on the sheet of $\sqrt{R(z')} \equiv \sqrt{(z' - \alpha)(z' - \alpha')(z' - \beta')(z' - \beta)}$ for which $\sqrt{R(z')} > 0$ for $z' > \beta$, with cuts on $[\alpha, \alpha']$ and $[\beta', \beta]$.

For a proof of this proposition see the next section.

2.4. **g -function.** Define the g -function on $\mathbb{C} \setminus [-\infty, \beta]$ as

$$g(z) = \int_{\alpha}^{\beta} \log(z - x) d\nu_0(x) \quad (2.22)$$

where we take the principal branch for logarithm.

Properties of $g(z)$:

- (1) $g(z)$ is analytic in $\mathbb{C} \setminus (-\infty, \beta]$.
- (2) For large z ,

$$g(z) = \log z - \sum_{j=1}^{\infty} \frac{g_j}{z^j}, \quad g_j = \int_{\alpha}^{\beta} \frac{x^j}{j} d\nu_0(x). \quad (2.23)$$

(3)

$$g'(z) = \omega(z). \quad (2.24)$$

(4) From the first relation in (2.12) we have that

$$g_+(x) + g_-(x) = |x| - \zeta x + l \quad \text{for } x \in [\alpha, \alpha'] \cup [\beta', \beta], \quad (2.25)$$

where g_+ and g_- refer to the limiting values of g from the upper and lower half-planes, respectively. By differentiating this equation we obtain that

$$\omega_+(x) + \omega_-(x) = g'_+(x) + g'_-(x) = \operatorname{sgn} x - \zeta \quad \text{for } x \in [\alpha, \alpha'] \cup [\beta', \beta], \quad (2.26)$$

Consider the function

$$f(x) = g_+(x) + g_-(x) - (|x| - \zeta x + l). \quad (2.27)$$

We have from (2.25), (2.26) that

$$f(x) = f'(x) = 0 \quad \text{for } x = \alpha, \alpha', \beta', \beta, \quad (2.28)$$

and from (2.21) that

$$f''(x) = -\frac{1}{\sqrt{(x - \alpha)(x - \alpha')(x - \beta')(x - \beta)}} \quad \text{for } x \in (-\infty, \alpha) \cup (\alpha', \beta') \cup (\beta, \infty). \quad (2.29)$$

Since

$$f''(x) < 0 \quad \text{for } x \in (-\infty, \alpha) \cup (\beta, \infty). \quad (2.30)$$

and

$$f''(x) > 0 \quad \text{for } x \in (\alpha', \beta'), \quad x \neq 0, \quad (2.31)$$

we obtain that

$$g_+(x) + g_-(x) \begin{cases} = |x| - \zeta x + l & \text{for } x \in [\alpha, \alpha'] \cup [\beta', \beta], \\ > |x| - \zeta x + l & \text{for } x \in (\alpha', \beta'), \\ < |x| - \zeta x + l & \text{for } x \in \mathbb{R} \setminus [\alpha, \beta]. \end{cases} \quad (2.32)$$

(5) Equation (2.22) implies that the function

$$G(x) \equiv g_+(x) - g_-(x) \quad (2.33)$$

is pure imaginary for all real x , and

$$G(x) = \begin{cases} 2\pi i & \text{for } -\infty < x \leq \alpha, \\ 2\pi i - 2\pi i \int_{\alpha}^x \rho(s) ds & \text{for } \alpha \leq x \leq \alpha', \\ 2\pi i \left(\frac{1+\zeta}{2} - \frac{x}{2\gamma} \right) & \text{for } \alpha' \leq x \leq \beta', \\ 2\pi i \int_x^{\beta} \rho(s) ds & \text{for } \beta' \leq x \leq \beta, \\ 0 & \text{for } \beta \leq x < \infty. \end{cases} \quad (2.34)$$

From (2.32) and (2.34) we obtain that

$$2g_{\pm}(x) = \begin{cases} |x| - \zeta x + l \pm \left[2\pi i - 2\pi i \int_{\alpha}^x \rho(s) ds \right] & \text{for } \alpha \leq x \leq \alpha', \\ |x| - \zeta x + l \pm 2\pi i \int_x^{\beta} \rho(s) ds & \text{for } \beta' \leq x \leq \beta. \end{cases} \quad (2.35)$$

(6) Also, from (2.34)

$$\left. \frac{dG(x+iy)}{dy} \right|_{y=0} = 2\pi\rho(x) > 0, \quad x \in (\alpha, \beta). \quad (2.36)$$

Observe that from (2.25) we have that

$$G(x) = 2g_+(x) - V(x) - l = -[2g_-(x) - V(x) - l], \quad x \in [\alpha, \alpha'] \cup [\beta', \beta], \quad (2.37)$$

where $V(x) \equiv |x| - \zeta x$.

2.5. Evaluation of the Lagrange multiplier l . We have the following proposition.

Proposition 2.3. *The Lagrange multiplier l solves the equation,*

$$e^{\frac{l}{2}} = \frac{\pi \vartheta_1'(0)}{2e\vartheta_1(\omega)}. \quad (2.38)$$

For a proof of this proposition see the next section.

3. PROOF OF PROPOSITIONS 2.1, 2.2, AND 2.3

Proof of Proposition 2.1. Following Zinn-Justin [31], we make the following elliptic change of variables:

$$u(z) = \frac{1}{2} \sqrt{(\beta' - \alpha)(\beta - \alpha')} \int_{\beta}^z \frac{dz'}{\sqrt{(z' - \alpha)(z' - \alpha')(z' - \beta')(z' - \beta)}}, \quad (3.1)$$

where integration takes place on the sheet on $\sqrt{R(z')}$ specified in Proposition 2.2. To understand this integral in terms of the usual elliptic integrals, we first make the change of variables

$$v(z') = \frac{(\beta - z')(\beta' - \alpha)}{(\beta' - z')(\beta - \alpha)}, \quad (3.2)$$

so that

$$z' = \frac{\beta'(\beta - \alpha)v - \beta(\beta' - \alpha)}{(\beta - \alpha)v - (\beta' - \alpha)}. \quad (3.3)$$

Note that $v(\beta) = 0$, $v(\beta') = \infty$, and $v(\alpha) = 1$. When we substitute v into equation (3.1), we have

$$u(z) = \frac{1}{2k} \int_0^{v(z)} \frac{dv}{\sqrt{v(v-1)(v-\frac{1}{k^2})}}, \quad (3.4)$$

where

$$k = \sqrt{\frac{(\beta - \alpha)(\beta' - \alpha')}{(\beta' - \alpha)(\beta - \alpha')}}. \quad (3.5)$$

We next take $v' = \sqrt{v}$, obtaining

$$u(z) = \int_0^{\sqrt{v(z)}} \frac{dv'}{\sqrt{(1-v'^2)(1-k^2v'^2)}}, \quad (3.6)$$

which corresponds to $\sqrt{v(z)} = \text{sn}(u, k)$, so that

$$\frac{(\beta - z)(\beta' - \alpha)}{(\beta' - z)(\beta - \alpha)} = \text{sn}^2(u), \quad \text{sn}(u) = \text{sn}(u, k). \quad (3.7)$$

Notice that u maps the upper z -plane conformally and bijectively onto the rectangle $[0, K] \times [0, iK']$, and the lower z -plane conformally and bijectively onto the rectangle $[0, K] \times [-iK', 0]$, where

$$\begin{aligned} K = u(\alpha) &= \int_0^1 \frac{dv'}{\sqrt{(1-v'^2)(1-k^2v'^2)}} \quad \text{and} \\ K' = -iu(\beta') &= \int_1^{\frac{1}{k}} \frac{dv'}{\sqrt{(v'^2-1)(1-k^2v'^2)}} \end{aligned} \quad (3.8)$$

are the usual complete integrals of the first kind. More specifically (see Fig. 6),

- (1) The upper (resp. lower) cusp of the interval $[\beta', \beta]$ is mapped onto the interval $[0, iK']$ (resp. $[0, -iK']$).
- (2) The upper (resp. lower) cusp of the interval $[\alpha, \alpha']$ is mapped onto the interval $[K, K + iK']$ (resp. $[K, K - iK']$).
- (3) The interval $[\alpha', \beta']$ is mapped onto the interval $[iK', K + iK']$ or the interval $[-iK', K - iK']$, depending on the path of integration.
- (4) The remaining part of the real axis, $[-\infty, \alpha] \cup [\beta, \infty]$, is mapped onto the interval $[0, K]$, with $u(\infty) = u_* = u_\infty$.

We will denote the rectangle $[0, K] \times [-iK', iK']$ as R , the *fundamental domain* of the function $z(u)$. We can now define

$$\tilde{\omega}(u) = \omega(z(u)) \quad \text{for } u \in R. \quad (3.9)$$




FIGURE 6. The u -plane. Here $u_* = u_\infty \equiv u(\infty)$ and $a = u_* + iK'$.

The Euler-Lagrange equation (2.13) and the equation (2.14) then become

$$\begin{aligned}
 \tilde{\omega}(u) + \tilde{\omega}(-u) &= 1 - \zeta & \text{for } u \in [-iK', iK'] \\
 \tilde{\omega}(u) + \tilde{\omega}(-u + 2K) &= -1 - \zeta & \text{for } u \in [K - iK', K + iK'] \\
 \tilde{\omega}(u + 2iK') - \tilde{\omega}(u) &= -\frac{i\pi}{\gamma} & \text{for } u \in [-iK', K - iK'].
 \end{aligned} \tag{3.10}$$

The function $\omega(z)$ is analytic in $\mathbb{C} \setminus [\alpha, \beta]$, but can be analytically continued from either above or below through any of the cuts $[\alpha, \alpha']$, $[\alpha', \beta']$, and $[\beta', \beta]$. These analytic continuations in the z -plane give an analytic continuation of $\tilde{\omega}$ in the u -plane into a neighborhood of R , which can then be continued by equations (3.10) to the entire u -plane. We therefore have that $\tilde{\omega}$ is analytic and satisfies equations (3.10) throughout the u -plane. The first two equations of (3.10) can be combined as

$$\tilde{\omega}(u + 2K) = \tilde{\omega}(u) - 2. \tag{3.11}$$

It therefore follows that $\tilde{\omega}$ is a linear function of u , as its derivative is a doubly periodic entire function. We also know from the fact that $\omega(z) \sim \frac{1}{z}$ at infinity that

$$\tilde{\omega}(u) = -\frac{2}{\sqrt{(\beta' - \alpha)(\beta - \alpha')}}(u - u_\infty) + O(u - u_\infty)^2 \tag{3.12}$$

in some neighborhood of u_∞ , where u_∞ is the image of infinity under the map $u(z)$. It thus follows from (3.10), (3.11), and (3.12) that

$$\tilde{\omega}(u) = -\frac{1}{K}(u - u_\infty) \tag{3.13}$$

and that

$$\frac{K'}{K} = \frac{\pi}{2\gamma} \quad (3.14)$$

$$\sqrt{(\beta' - \alpha)(\beta - \alpha')} = 2K \quad (3.15)$$

$$\frac{u_\infty}{K} = \frac{1 - \zeta}{2}. \quad (3.16)$$

From (3.7) we obtain that

$$\frac{\beta' - \alpha}{\beta - \alpha} = \text{sn}^2(u_\infty). \quad (3.17)$$

This implies that

$$\begin{aligned} \text{cn}^2(u_\infty) &= 1 - \text{sn}^2(u_\infty) = 1 - \frac{\beta' - \alpha}{\beta - \alpha} = \frac{\beta - \beta'}{\beta - \alpha}, \\ \text{dn}^2(u_\infty) &= 1 - k^2 \text{sn}^2(u_\infty) = 1 - \frac{(\beta - \alpha)(\beta' - \alpha')}{(\beta' - \alpha)(\beta - \alpha')} \frac{(\beta' - \alpha)}{(\beta - \alpha)} = \frac{\beta - \beta'}{\beta - \alpha'}. \end{aligned} \quad (3.18)$$

From equations (3.15), (3.17), (3.18) we obtain the distances between the turning points in terms of u_∞ :

$$\begin{aligned} \beta - \alpha &= 2K \frac{\text{dn}(u_\infty)}{\text{sn}(u_\infty) \text{cn}(u_\infty)} \\ \beta - \alpha' &= 2K \frac{\text{cn}(u_\infty)}{\text{sn}(u_\infty) \text{dn}(u_\infty)} \\ \beta - \beta' &= 2K \frac{\text{cn}(u_\infty) \text{dn}(u_\infty)}{\text{sn}(u_\infty)}. \end{aligned} \quad (3.19)$$

The functions sn, cn, and dn are expressed in terms of Jacobi theta functions as follows (see, eg. [30]),

$$\text{sn}(u) = \frac{\vartheta_3(0) \vartheta_1(\frac{\pi u}{2K})}{\vartheta_2(0) \vartheta_4(\frac{\pi u}{2K})}, \quad \text{cn}(u) = \frac{\vartheta_4(0) \vartheta_2(\frac{\pi u}{2K})}{\vartheta_2(0) \vartheta_4(\frac{\pi u}{2K})}, \quad \text{dn}(u) = \frac{\vartheta_4(0) \vartheta_3(\frac{\pi u}{2K})}{\vartheta_3(0) \vartheta_4(\frac{\pi u}{2K})}. \quad (3.20)$$

By (3.14), the half-period ratio τ and the elliptic nome q of the theta functions are

$$\tau = \frac{iK'}{K} = \frac{i\pi}{2\gamma} \quad \text{and} \quad q = e^{\frac{-\pi K'}{K}} = e^{\frac{-\pi^2}{2\gamma}}. \quad (3.21)$$

If we take into account the fact that

$$\vartheta_3(0)^2 = \frac{2K}{\pi} \quad (3.22)$$

and equation (3.16), we can write equations for the distances between the turning points that involve the original parameters only:

$$\begin{aligned} \beta - \alpha &= \pi \vartheta_2^2(0) \frac{\vartheta_3(\frac{\omega}{2}) \vartheta_4(\frac{\omega}{2})}{\vartheta_1(\frac{\omega}{2}) \vartheta_2(\frac{\omega}{2})}, & \beta - \alpha' &= \pi \vartheta_3(0)^2 \frac{\vartheta_1(\frac{\omega}{2}) \vartheta_3(\frac{\omega}{2})}{\vartheta_2(\frac{\omega}{2}) \vartheta_4(\frac{\omega}{2})}, \\ \beta' - \alpha &= \pi \vartheta_3^2(0) \frac{\vartheta_2(\frac{\omega}{2}) \vartheta_4(\frac{\omega}{2})}{\vartheta_1(\frac{\omega}{2}) \vartheta_3(\frac{\omega}{2})}, \end{aligned} \quad (3.23)$$

giving (2.17). These equations determine the end-points $\alpha, \alpha', \beta', \beta$ up to a shift. To fix the shift we use the equation (2.12) at the points α' and β' to obtain

$$\int_{\alpha'}^{\beta'} (\omega(z + i0) + \omega(z - i0)) dz = (1 - \zeta)\beta' + (1 + \zeta)\alpha'. \quad (3.24)$$

Writing this integral in terms of u gives

$$\int_{iK'}^{K+iK'} \frac{1}{K} (u - u_\infty) r'(u) du + \int_{-iK'}^{K-iK'} \frac{1}{K} (u - u_\infty) r'(u) du = (1 - \zeta)\beta' + (1 + \zeta)\alpha' \quad (3.25)$$

where

$$r(u) = \frac{\beta'(\beta - \alpha)\text{sn}^2(u) - \beta(\beta' - \alpha)}{(\beta - \alpha)\text{sn}^2(u) - (\beta' - \alpha)} = \frac{\beta - \frac{\beta'\text{sn}^2(u)}{\text{sn}^2(u_\infty)}}{1 - \frac{\text{sn}^2(u)}{\text{sn}^2(u_\infty)}} \quad \text{and} \quad r'(u) = \frac{d}{du} r(u). \quad (3.26)$$

Note that $r(\pm iK') = \beta'$ and $r(K \pm iK') = \alpha'$. Integrating by parts gives

$$\frac{2}{K} ((K - u_\infty)\alpha' + \beta' u_\infty) - \int_{iK'}^{K+iK'} \frac{r(u)}{K} du - \int_{-iK'}^{K-iK'} \frac{r(u)}{K} du = (1 - \zeta)\beta' + (1 + \zeta)\alpha' \quad (3.27)$$

or equivalently

$$\int_{iK'}^{K+iK'} r(u) du + \int_{-iK'}^{K-iK'} r(u) du = 0. \quad (3.28)$$

We can evaluate these integrals by first writing $r(u)$ in the form

$$r(u) = \beta + \left(\frac{\beta - \beta'}{\text{sn}^2(u_\infty)} \right) \frac{\text{sn}^2(u)}{1 - \frac{\text{sn}^2(u)}{\text{sn}^2(u_\infty)}}. \quad (3.29)$$

and using the functions

$$\Theta(u) = \vartheta_4\left(\frac{\pi u}{2K}\right), \quad Z(u) = \frac{\Theta'(u)}{\Theta(u)}. \quad (3.30)$$

The addition formulae for the sn and Z functions are (see [30]):

$$\begin{aligned} \text{sn}(u \pm a) &= \frac{\text{sn}(u)\text{cn}(a)\text{dn}(a) \pm \text{sn}(a)\text{cn}(u)\text{dn}(u)}{1 - k^2\text{sn}^2(a)\text{sn}^2(u)}, \\ Z(u \pm a) &= Z(u) \pm Z(a) \mp k^2\text{sn}(u)\text{sn}(a)\text{sn}(u \pm a). \end{aligned} \quad (3.31)$$

Thus we have

$$\begin{aligned} Z(u - a) - Z(u + a) + 2Z(a) &= k^2\text{sn}(u)\text{sn}(a)(\text{sn}(u + a) + \text{sn}(u - a)) \\ &= \frac{k^2\text{sn}(u)\text{sn}(a) [2\text{sn}(u)\text{cn}(a)\text{dn}(a)]}{1 - k^2\text{sn}^2(a)\text{sn}^2(u)} \\ &= \frac{2k^2\text{sn}(a)\text{cn}(a)\text{dn}(a)\text{sn}^2(u)}{1 - k^2\text{sn}^2(a)\text{sn}^2(u)}. \end{aligned} \quad (3.32)$$

We also have the half- and quarter-period identities

$$\text{sn}(u + iK') = \frac{1}{k\text{sn}(u)}, \quad \text{cn}(u + iK') = \frac{-i \text{dn}(u)}{k \text{sn}(u)}, \quad \text{dn}(u + iK') = -i \frac{\text{cn}(u)}{\text{sn}(u)}. \quad (3.33)$$

In particular, notice that $\frac{1}{\text{sn}(u_\infty)} = k\text{sn}(u_\infty + iK')$. Using the addition formulae (3.32) we can write $r(u)$ as

$$r(u) = \beta + \left(\frac{\beta - \beta'}{2k^2\text{sn}(a)\text{cn}(a)\text{dn}(a)\text{sn}^2(u_\infty)} \right) (Z(u - a) - Z(u + a) + 2Z(a)), \quad (3.34)$$

where $a = u_\infty + iK'$, (see Fig. 6). From (3.33) and (3.19), it follows that

$$\frac{\beta - \beta'}{2k^2\text{sn}(u_\infty + iK')\text{cn}(u_\infty + iK')\text{dn}(u_\infty + iK')\text{sn}^2(u_\infty)} = -K. \quad (3.35)$$

Thus we can write (3.34) as

$$r(u) = \beta - K [Z(u - u_\infty - iK') - Z(u + u_\infty + iK') + 2Z(u_\infty + iK')]. \quad (3.36)$$

If we write $u = x + iK'$ in the first integral of (3.28), and $u = x - iK'$ in the second, we obtain

$$\int_0^K [2\beta - 4KZ(u_\infty + iK') - K[Z(x - u_\infty) - Z(x + u_\infty + 2iK') + Z(x - u_\infty - 2iK') - Z(x + u_\infty)]] dx = 0 \quad (3.37)$$

From the periodic properties of ϑ_4 , it follows that

$$Z(u \pm 2iK') = Z(u) \mp \frac{\pi i}{K}, \quad (3.38)$$

so we can write (3.37) as

$$\int_0^K \left(2\beta - 4KZ(u_\infty + iK') - 2\pi i + 2K[Z(x + u_\infty) - Z(x - u_\infty)] \right) dx = 0 \quad (3.39)$$

This equation is readily integrated, as Z is the logarithmic derivative of the Θ function. Integrating gives

$$\begin{aligned} 0 &= \left[(2\beta - 4KZ(u_\infty + iK') - 2\pi i)x + 2K \log \frac{\Theta(x + u_\infty)}{\Theta(x - u_\infty)} \right]_{x=0}^K \\ &= 2K\beta - 4K^2Z(u_\infty + iK') - 2K\pi i + 2K \log \left(\frac{\Theta(K + u_\infty)}{\Theta(K - u_\infty)} \frac{\Theta(-u_\infty)}{\Theta(u_\infty)} \right). \end{aligned} \quad (3.40)$$

The logarithmic term in this equation is zero due to the evenness and periodicity (period $2K$) of the Θ function and the fact that the relevant term in the integration is real on the entire contour of integration. Thus we have that

$$\beta = 2KZ(u_\infty + iK') + \pi i. \quad (3.41)$$

From (1.42), we can deduce that

$$Z(u_\infty + iK') = -\frac{\pi}{2K} \left(\frac{\vartheta'_2(\frac{\omega}{2})}{\vartheta_2(\frac{\omega}{2})} + i \right) \quad (3.42)$$

and write (3.41) as

$$\beta = -\pi \frac{\vartheta'_2(\frac{\omega}{2})}{\vartheta_2(\frac{\omega}{2})}. \quad (3.43)$$

This equation, together with equations (3.23) and (3.21), determine the end-points, $\alpha, \alpha', \beta', \beta$. In fact, similar to (3.43) we have the following explicit formulae for the other end-points:

$$\alpha = -\pi \frac{\vartheta'_1(\frac{\omega}{2})}{\vartheta_1(\frac{\omega}{2})}, \quad \alpha' = -\pi \frac{\vartheta'_4(\frac{\omega}{2})}{\vartheta_4(\frac{\omega}{2})}, \quad \beta' = -\pi \frac{\vartheta'_3(\frac{\omega}{2})}{\vartheta_3(\frac{\omega}{2})}. \quad (3.44)$$

This follows from (3.23), (3.43), and the identities (E.2). Similarly, in addition to the formulae (3.23) for distances between turning points, we get (2.16).

Proof of the formula of Zinn-Justin for $\frac{\alpha+\alpha'+\beta'+\beta}{4}$. We immediately have from (3.44) that

$$\frac{\alpha + \alpha' + \beta' + \beta}{4} = -\frac{\pi}{4} \left[\frac{\vartheta'_1(\frac{\omega}{2})}{\vartheta_1(\frac{\omega}{2})} + \frac{\vartheta'_2(\frac{\omega}{2})}{\vartheta_2(\frac{\omega}{2})} + \frac{\vartheta'_3(\frac{\omega}{2})}{\vartheta_3(\frac{\omega}{2})} + \frac{\vartheta'_4(\frac{\omega}{2})}{\vartheta_4(\frac{\omega}{2})} \right]. \quad (3.45)$$

From the identity (E.6), we can deduce that

$$\frac{\vartheta'_1(2z)}{\vartheta_1(2z)} = \frac{1}{2} \left[\frac{\vartheta'_1(z)}{\vartheta_1(z)} + \frac{\vartheta'_2(z)}{\vartheta_2(z)} + \frac{\vartheta'_3(z)}{\vartheta_3(z)} + \frac{\vartheta'_4(z)}{\vartheta_4(z)} \right], \quad (3.46)$$

thus we have

$$\frac{\alpha + \alpha' + \beta' + \beta}{4} = -\frac{\pi}{2} \frac{\vartheta'_1(\omega)}{\vartheta_1(\omega)} = -\frac{\pi}{2} \frac{\vartheta'_1(\frac{\pi}{2} + \frac{\pi\zeta}{2})}{\vartheta_1(\frac{\pi}{2} + \frac{\pi\zeta}{2})} = -\frac{\pi}{2} \frac{\vartheta'_2(\frac{\pi\zeta}{2})}{\vartheta_2(\frac{\pi\zeta}{2})}. \quad (3.47)$$

Proof of Proposition 2.2. From equations (3.13), (3.1), and (3.15) we obtain formula (2.21), cf. [31]. From formula (2.21) and equations (2.8), (2.14) we obtain that the equilibrium density function $\rho(x)$ is given by formulae (2.19). We are left to prove formula (2.20).

By (3.1), (3.15), and (2.19), on the interval $[\beta', \beta]$,

$$\rho(x) = \frac{1}{iK\pi} u_+(x) \quad \text{for } x \in [\beta', \beta]. \quad (3.48)$$

It follows that

$$\int_{\beta'}^{\beta} \rho(x) dx = \frac{1}{iK\pi} \int_{\beta'}^{\beta} u_+(x) dx = \frac{1}{iK\pi} \int_{iK'}^0 ur'(u) du, \quad (3.49)$$

where $r(u)$ is defined in (3.26). If we use equation (3.36), together with formula (3.41), we can write $r(u)$ as

$$r(u) = i\pi - K[Z(u - u_\infty - iK') - Z(u + u_\infty + iK')] \quad (3.50)$$

Integrating (3.49) by parts, we get

$$\int_{iK'}^0 ur'(u) du = -\beta' iK' - \pi K' - K \left[\log \left(\frac{\Theta(-u_\infty - iK')\Theta(u_\infty + 2iK')}{\Theta(-u_\infty)\Theta(u_\infty + iK')} \right) \right]. \quad (3.51)$$

Using the fact that Θ is an even function and the identity

$$\Theta(u + 2iK') = e^{i\pi} e^{-i\pi\tau} e^{\frac{-i\pi u}{K}} \Theta(u), \quad (3.52)$$

we can write (3.51) as

$$\begin{aligned} \int_{iK'}^0 ur'(u) du &= -\beta' iK' - \pi K' + K \left(i\pi + \pi \frac{K'}{K} - \frac{i\pi u_\infty}{K} \right) \\ &= i(K\pi - \beta' K' - \pi u_\infty). \end{aligned} \quad (3.53)$$

Remark: There is some question here as to which branch of the logarithm to take, but it is clear that we have chosen the correct branch, as it is the only one that gives $0 < \int_{\beta'}^{\beta} \rho(x) dx < 1$.

Thus, from (3.49) and (3.53), we have

$$\int_{\beta'}^{\beta} \rho(x) dx = \frac{1}{iK\pi} i(K\pi - \beta'K' - \pi u_{\infty}) = 1 - \frac{\beta'K'}{\pi K} - \frac{u_{\infty}}{K} = 1 - \frac{\beta'}{2\gamma} - \frac{1-\zeta}{2}, \quad (3.54)$$

hence by (2.19),

$$\int_0^{\beta} \rho(x) dx = \int_0^{\beta'} \rho(x) dx + \int_{\beta'}^{\beta} \rho(x) dx = \frac{\beta'}{2\gamma} + 1 - \frac{\beta'}{2\gamma} - \frac{1-\zeta}{2} = \frac{1+\zeta}{2}, \quad (3.55)$$

which proves formula (2.20).

Proof of Proposition 2.3. By taking $x = \beta$ we obtain from (2.32) that

$$l = 2g(\beta) - V(\beta) = 2g(\beta) - (1-\zeta)\beta. \quad (3.56)$$

We also have that

$$\lim_{A \rightarrow \infty} [g(A) - \log A] = 0, \quad (3.57)$$

hence

$$\begin{aligned} l &= -2 \lim_{A \rightarrow \infty} [g(A) - g(\beta) - \log A] - (1-\zeta)\beta \\ &= -2 \lim_{A \rightarrow \infty} \left[\int_{\beta}^A \omega(z) dz - \log A \right] - (1-\zeta)\beta. \end{aligned} \quad (3.58)$$

Writing this integral in terms of u (so $z = r(u)$) gives

$$\begin{aligned} l &= 2 \lim_{A \rightarrow \infty} \left[\int_0^B \frac{1}{K} (u - u_{\infty}) r'(u) du + \log A \right] - (1-\zeta)\beta \\ &= 2 \lim_{B \rightarrow u_{\infty}} \left[\int_0^B \frac{1}{K} (u - u_{\infty}) r'(u) du + \log r(B) \right] - (1-\zeta)\beta, \end{aligned} \quad (3.59)$$

where $A = r(B)$. Integrating by parts gives

$$l = 2 \lim_{B \rightarrow u_{\infty}} \left[\frac{1}{K} (u - u_{\infty}) r(u) \Big|_{u=0}^B - \frac{1}{K} \int_0^B r(u) du + \log r(B) \right] - (1-\zeta)\beta. \quad (3.60)$$

From (3.26) we obtain that $r(0) = \beta$ and

$$\lim_{B \rightarrow u_{\infty}} (B - u_{\infty}) r(B) = -\frac{(\beta - \beta') \operatorname{sn}(u_{\infty})}{2 \operatorname{sn}'(u_{\infty})} = -\frac{(\beta - \beta') \operatorname{sn}(u_{\infty})}{2 \operatorname{cn}(u_{\infty}) \operatorname{dn}(u_{\infty})} = -K, \quad (3.61)$$

hence

$$\begin{aligned} l &= 2 \left[-1 + \frac{\beta(1-\zeta)}{2} \right] - 2 \lim_{B \rightarrow u_{\infty}} \left[\frac{1}{K} \int_0^B r(u) du - \log r(B) \right] - (1-\zeta)\beta \\ &= -2 - 2 \lim_{B \rightarrow u_{\infty}} \left[\frac{1}{K} \int_0^B r(u) du - \log r(B) \right]. \end{aligned} \quad (3.62)$$

Using equation (3.50) for $r(u)$, we immediately get that

$$\frac{1}{K} \int_0^B r(u) du = \frac{B\pi i}{K} + \log \left[\frac{\Theta(B + u_\infty + iK')}{\Theta(B - u_\infty - iK')} \right] \quad (3.63)$$

Now using equation (3.26) for $r(u)$, we have

$$\begin{aligned} & \lim_{B \rightarrow u_\infty} \left[\frac{1}{K} \int_0^B r(u) du - \log r(B) \right] \\ &= \frac{u_\infty \pi i}{K} + \lim_{B \rightarrow u_\infty} \log \left[\frac{\Theta(B + u_\infty + iK')(\operatorname{sn}^2(u_\infty) - \operatorname{sn}^2(B))}{\Theta(B - u_\infty - iK')(\beta \operatorname{sn}^2(u_\infty) - \beta' \operatorname{sn}^2(B))} \right] \\ &= \frac{u_\infty \pi i}{K} + \log \left[\frac{\Theta(2u_\infty + iK') 2 \operatorname{sn}(u_\infty) \operatorname{sn}'(u_\infty)}{\Theta'(iK')(\beta - \beta') \operatorname{sn}^2(u_\infty)} \right] \\ &= \frac{u_\infty \pi i}{K} + \log \left[\frac{2e^{\frac{-i\pi u_\infty}{K}} \vartheta_1\left(\frac{\pi u_\infty}{K}\right)}{\pi \vartheta_1'(0)} \right] \\ &= \log \left[\frac{2\vartheta_1(\omega)}{\pi \vartheta_1'(0)} \right]. \end{aligned} \quad (3.64)$$

Plugging this into (3.62) gives

$$l = -2 + 2 \log \left(\frac{\pi \vartheta_1'(0)}{2\vartheta_1(\omega)} \right), \quad (3.65)$$

and thus we obtain that

$$e^{\frac{l}{2}} = \frac{\pi \vartheta_1'(0)}{2e\vartheta_1(\omega)}. \quad (3.66)$$

4. RIEMANN-HILBERT APPROACH: INTERPOLATION PROBLEM

The Riemann-Hilbert approach to discrete orthogonal polynomials is based on the following Interpolation Problem (IP), which was introduced in the paper [9] of Borodin and Borovichenko under the name of the discrete Riemann-Hilbert problem. See also the monograph [5] of Baik, Kriecherbauer, McLaughlin, and Miller, in which it is called the Interpolation Problem.

We will consider the lattice L_n defined in (1.27) and the weight $w_n(x)$ defined in (1.25).

Interpolation Problem. For a given $n = 0, 1, \dots$, find a 2×2 matrix-valued function $\mathbf{P}_n(z) = (\mathbf{P}_{nij}(z))_{1 \leq i, j \leq 2}$ with the following properties:

- (1) *Analyticity:* $\mathbf{P}_n(z)$ is an analytic function of z for $z \in \mathbb{C} \setminus L_n$.
- (2) *Residues at poles:* At each node $x \in L_n$, the elements $\mathbf{P}_{n11}(z)$ and $\mathbf{P}_{n21}(z)$ of the matrix $\mathbf{P}_n(z)$ are analytic functions of z , and the elements $\mathbf{P}_{n12}(z)$ and $\mathbf{P}_{n22}(z)$ have a simple pole with the residues,

$$\operatorname{Res}_{z=x} \mathbf{P}_{nj2}(z) = w_n(x) \mathbf{P}_{nj1}(x), \quad j = 1, 2. \quad (4.1)$$

- (3) *Asymptotics at infinity:* There exists a function $r(x) > 0$ on L_n such that

$$\lim_{x \rightarrow \infty} r(x) = 0, \quad (4.2)$$

and such that as $z \rightarrow \infty$, $\mathbf{P}_n(z)$ admits the asymptotic expansion,

$$\mathbf{P}_n(z) \sim \left(\mathbf{I} + \frac{\mathbf{P}_1}{z} + \frac{\mathbf{P}_2}{z^2} + \dots \right) \begin{pmatrix} z^n & 0 \\ 0 & z^{-n} \end{pmatrix}, \quad z \in \mathbb{C} \setminus \left[\bigcup_{x \in L_n}^\infty D(x, r(x)) \right], \quad (4.3)$$

where $D(x, r(x))$ denotes a disk of radius $r(x) > 0$ centered at x and $\mathbf{I}$ is the identity matrix.

It is not difficult to see (see [9] and [5]) that the IP has a unique solution, which is

$$\mathbf{P}_n(z) = \begin{pmatrix} P_{nn}(z) & C(w_n P_{nn})(z) \\ (h_{n,n-1})^{-1} P_{n,n-1}(z) & (h_{n,n-1})^{-1} C(w_n P_{n,n-1})(z) \end{pmatrix} \quad (4.4)$$

where the Cauchy transformation C is defined by the formula,

$$C(f)(z) = \sum_{x \in L_n} \frac{f(x)}{z - x}, \quad (4.5)$$

and $P_{nk}(z) = z^k + \dots$ are monic polynomials orthogonal with the weight $w_n(x)$. Because of the orthogonality condition, as $z \rightarrow \infty$,

$$C(w_n P_{nk})(z) = \sum_{x \in L_n} \frac{w_n(x) P_{nk}(x)}{z - x} \sim \sum_{x \in L_n} w_n(x) P_{nk}(x) \sum_{j=0}^{\infty} \frac{x^j}{z^{j+1}} = \frac{h_{nk}}{z^{k+1}} + \sum_{j=k+2}^{\infty} \frac{a_j}{z^j}, \quad (4.6)$$

which justifies asymptotic expansion (4.3). We have that

$$h_{nn} = [\mathbf{P}_1]_{12}, \quad h_{n,n-1}^{-1} = [\mathbf{P}_1]_{21}. \quad (4.7)$$

5. REDUCTION OF IP TO RHP

5.1. Preliminary Considerations. We would like to reduce the Interpolation Problem to a Riemann-Hilbert Problem (RHP). Introduce the function,

$$\Pi(z) = \frac{2\gamma}{n\pi} \sin \frac{n\pi z}{2\gamma}. \quad (5.1)$$

Observe that

$$\Pi(x_k) = 0, \quad \Pi'(x_k) = (-1)^k, \quad \exp\left(\frac{in\pi x_k}{2\gamma}\right) = (-1)^k \quad \text{for } x_k = \frac{2\gamma k}{n} \in L_n. \quad (5.2)$$

Introduce the upper triangular matrices,

$$\mathbf{D}_\pm^u(z) = \begin{pmatrix} 1 & -\frac{w_n(z)}{\Pi(z)} e^{\pm \frac{in\pi z}{2\gamma}} \\ 0 & 1 \end{pmatrix}, \quad (5.3)$$

and the lower triangular matrices,

$$\mathbf{D}_\pm^l = \begin{pmatrix} \Pi(z)^{-1} & 0 \\ -\frac{1}{w_n(z)} e^{\pm \frac{in\pi z}{2\gamma}} & \Pi(z) \end{pmatrix} = \begin{pmatrix} \Pi(z)^{-1} & 0 \\ 0 & \Pi(z) \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -\frac{1}{\Pi(z)w_n(z)} e^{\pm \frac{in\pi z}{2\gamma}} & 1 \end{pmatrix}. \quad (5.4)$$

Define the matrix-valued functions,

$$\mathbf{R}_n^u = \mathbf{P}_n(z) \times \begin{cases} \mathbf{D}_+^u(z), & \text{when } \operatorname{Im} z \geq 0, \\ \mathbf{D}_-^u(z), & \text{when } \operatorname{Im} z \leq 0, \end{cases} \quad (5.5)$$

and

$$\mathbf{R}_n^l = \mathbf{P}_n(z) \times \begin{cases} \mathbf{D}_+^l(z), & \text{when } \operatorname{Im} z \geq 0, \\ \mathbf{D}_-^l(z), & \text{when } \operatorname{Im} z \leq 0. \end{cases} \quad (5.6)$$

From (4.4) we have that

$$\mathbf{R}_n^u(z) = \begin{pmatrix} P_{nn}(z) & -\frac{w_n(z)P_{nn}(z)}{\Pi(z)}e^{\pm \frac{i n \pi z}{2\gamma}} + C(w_n P_{nn})(z) \\ h_{n,n-1}^{-1}P_{n,n-1}(z) & -\frac{w_n(z)h_{n,n-1}^{-1}P_{n,n-1}(z)}{\Pi(z)}e^{\pm \frac{i n \pi z}{2\gamma}} + h_{n,n-1}^{-1}C(w_n P_{n,n-1})(z) \end{pmatrix}, \quad (5.7)$$

when $\pm \operatorname{Im} z \geq 0$,

and

$$\mathbf{R}_n^l(z) = \begin{pmatrix} \frac{P_{nn}(z)}{\Pi(z)} - \frac{C(w_n P_{nn})(z)}{w_n(z)}e^{\pm \frac{i n \pi z}{2\gamma}} & \Pi(z)C(w_n P_{nn})(z) \\ \frac{h_{n,n-1}^{-1}P_{n,n-1}(z)}{\Pi(z)} - \frac{h_{n,n-1}^{-1}C(w_n P_{n,n-1})(z)}{w_n(z)}e^{\pm \frac{i n \pi z}{2\gamma}} & \Pi(z)h_{n,n-1}^{-1}C(w_n P_{n,n-1})(z) \end{pmatrix}, \quad (5.8)$$

when $\pm \operatorname{Im} z \geq 0$.

Observe that the functions $\mathbf{R}_n^u(z)$, $\mathbf{R}_n^l(z)$ are meromorphic on the closed quadrants of the complex plane and they are two-valued on the real and imaginary axes. Their possible poles are located on the lattice L_n . An important result is that, in fact, due to some cancellations they do not have any poles at all. We have the following proposition.

Proposition 5.1. *The matrix-valued functions $\mathbf{R}_n^u(z)$ and $\mathbf{R}_n^l(z)$ have no poles and on the real line they satisfy the following jump conditions at $x \in \mathbb{R}$:*

$$\mathbf{R}_{n+}^u(x) = \mathbf{R}_{n-}^u(x)j_R^u(x), \quad j_R^u(x) = \begin{pmatrix} 1 & -\frac{n\pi i w_n(x)}{\gamma} \\ 0 & 1 \end{pmatrix}, \quad (5.9)$$

and

$$\mathbf{R}_{n+}^l(x) = \mathbf{R}_{n-}^l(x)j_R^l(x), \quad j_R^l(x) = \begin{pmatrix} 1 & 0 \\ -\frac{n\pi i}{\gamma w_n(x)} & 1 \end{pmatrix}, \quad (5.10)$$

Proof. It follows from the definition of $\mathbf{R}_n^u(z)$ that all possible poles of $\mathbf{R}_n^u(z)$ are located on the lattice L_n . Let us show that the residue of all these poles is equal to zero. Consider any $x_k \in L_n$. The residue of the matrix element $\mathbf{R}_{n,12}^u(z)$ at x_k is equal to

$$\operatorname{Res}_{z=x_k} \mathbf{R}_{n,12}^u(z) = -\frac{w_n(x_k)P_{nn}(x_k)}{(-1)^k}(-1)^k + w_n(x_k)P_{nn}(x_k) = 0. \quad (5.11)$$

Similarly we get that

$$\operatorname{Res}_{z=x_k} \mathbf{R}_{n,22}(z) = 0, \quad (5.12)$$

hence $\mathbf{R}_n^u(z)$ has no pole at x_k .

Similarly, the residue of the matrix element $\mathbf{R}_{n,11}^l(z)$ at x_k is equal to

$$\operatorname{Res}_{z=x_k} \mathbf{R}_{n,11}^l(z) = \frac{P_{nn}(x_k)}{(-1)^k} - \frac{w_n(x_k)P_{nn}(x_k)(-1)^k}{w_n(x_k)} = 0. \quad (5.13)$$

In the same way we obtain that

$$\operatorname{Res}_{z=x_k} \mathbf{R}_{n,21}(z) = 0. \quad (5.14)$$

In the entry $\mathbf{R}_{n,21}^l(z)$, the pole of the function $C(w_n P_n)(z)$ at $z = x_k$ is cancelled by the zero of the function $\Pi(z)$, hence $\mathbf{R}_{n,21}^l(z)$ has no pole at x_k . Similarly, $\mathbf{R}_{n,22}^l(z)$ has no pole at x_k as well, hence $\mathbf{R}_n^l(z)$ has no pole at x_k .

Let us evaluate to jump matrices at $x \in \mathbb{R}$. From (5.5) we have that

$$j_R^u(x) = \mathbf{D}_-^u(x)^{-1} \mathbf{D}_+^u(x) = \begin{pmatrix} 1 & -\frac{w_n(x)}{\Pi(x)} 2i \sin \frac{n\pi x}{2\gamma} \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & -\frac{n\pi i w_n(x)}{\gamma} \\ 0 & 1 \end{pmatrix}, \quad (5.15)$$

which proves (5.9). Similarly,

$$j_R^l(x) = \mathbf{D}_-^l(x)^{-1} \mathbf{D}_+^l(x) = \begin{pmatrix} 1 & 0 \\ -\frac{1}{\Pi(x)w_n(x)} 2i \sin \frac{n\pi x}{2\gamma} & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ -\frac{n\pi i}{\gamma w_n(x)} & 1 \end{pmatrix}, \quad (5.16)$$

which proves (5.10). $\square$

5.2. Reduction of IP to RHP. Let us discuss how to reduce the Interpolation Problem to a Riemann-Hilbert Problem. We follow the work [5] with some modifications. Denote

$$\Delta = L_n \cap [\alpha', \beta'], \quad \nabla = L_n \setminus \Delta. \quad (5.17)$$

Consider the oriented contour Σ on the complex plane depicted on Fig. 7, in which the horizontal lines are $\text{Im } z = \varepsilon, 0, -\varepsilon$, where $\varepsilon > 0$ is a small positive constant which will be determined later, and the vertical segments pass through the points $z = \alpha'$ and $z = \beta'$. Consider the regions $\Omega_{\pm}^{\Delta}$ and $\Omega_{\pm}^{\nabla}$ bounded by the contour Σ , see Fig. 7. Observe that the regions $\Omega_{\pm}^{\nabla}$ consist of two connected components, to the left and to the right of $\Omega_{\pm}^{\Delta}$.




FIGURE 7. The contour Σ .


Define

$$\mathbf{R}_n(z) = \begin{cases} \mathbf{K}_n \mathbf{R}_n^u(z) \mathbf{K}_n^{-1}, & \text{for } z \in \Omega_{\pm}^{\nabla}, \\ \mathbf{K}_n \mathbf{R}_n^l(z) \mathbf{K}_n^{-1}, & \text{for } z \in \Omega_{\pm}^{\Delta}, \\ \mathbf{K}_n \mathbf{P}_n(z) \mathbf{K}_n^{-1}, & \text{otherwise.} \end{cases} \quad (5.18)$$

where $\mathbf{K}_n = \begin{pmatrix} 1 & 0 \\ 0 & -\frac{n\pi i}{\gamma} \end{pmatrix}$.

Define a contour Σ_R by adding to the contour Σ a vertical segment $[i\varepsilon, -i\varepsilon]$, see Fig. 8. If $A \subset \mathbb{C}$ is a set on the complex plane and $b \in \mathbb{C}$ then, as usual, we denote

$$A + b = \{z = a + b, a \in A\}. \quad (5.19)$$

FIGURE 8. The contour Σ_R .

Proposition 5.2. *The matrix-valued function $\mathbf{R}_n(z)$ has the following jumps on the contour Σ_R :*

$$\mathbf{R}_{n+}(z) = \mathbf{R}_{n-}(z)j_R(z), \quad (5.20)$$

where

$$j_R(z) = \begin{cases} \begin{pmatrix} 1 & w(z) \\ 0 & 1 \end{pmatrix} & \text{when } z \in (-\infty, \alpha') \cup (\beta', \infty), \\ \begin{pmatrix} 1 & 0 \\ -(\frac{n\pi}{\gamma})^2 w(z)^{-1} & 1 \end{pmatrix} & \text{when } z \in [\alpha', \beta'], \\ \mathbf{K}_n \mathbf{D}_{\pm}^u(z) \mathbf{K}_n^{-1} = \begin{pmatrix} 1 & -\frac{i\gamma}{n\pi} \frac{w_n(z) e^{\pm \frac{in\pi z}{2\gamma}}}{\Pi(z)} \\ 0 & 1 \end{pmatrix} & \text{when } z \in (-\infty, \alpha') \cup (\beta', \infty) \pm i\varepsilon, \\ \mathbf{K}_n \mathbf{D}_{\pm}^l(z) \mathbf{K}_n^{-1} = \begin{pmatrix} \Pi(z)^{-1} & 0 \\ \frac{in\pi}{\gamma} \frac{e^{\pm \frac{in\pi z}{2\gamma}}}{w_n(z)} & \Pi(z) \end{pmatrix} & \text{when } z \in (\alpha', \beta') \pm i\varepsilon \\ \mathbf{K}_n \mathbf{D}_{\pm}^l(z)^{-1} \mathbf{D}_{\pm}^u(z) \mathbf{K}_n^{-1} = \begin{pmatrix} \Pi(z) & \frac{\gamma}{n\pi i} w_n(z) e^{\pm \frac{in\pi z}{2\gamma}} \\ -\frac{n\pi i}{\gamma} w_n(z)^{-1} e^{\pm \frac{in\pi z}{2\gamma}} & \mp \frac{n\pi i}{\gamma} e^{\pm \frac{in\pi z}{2\gamma}} \end{pmatrix} & \text{when } z \in (0, \pm i\varepsilon) + \alpha' \text{ or } z \in (0, \pm i\varepsilon) + \beta', \\ \mathbf{K}_n \mathbf{D}_{\pm}^0(z) \mathbf{K}_n^{-1} & \text{when } z \in (0, \pm i\varepsilon), \end{cases} \quad (5.21)$$

and

$$\mathbf{D}_{\pm}^0(z) = \begin{pmatrix} 1 & 0 \\ -\frac{2 \sinh(nz) e^{-n\zeta z} e^{\pm \frac{in\pi z}{2\gamma}}}{\Pi(z)} & 1 \end{pmatrix}. \quad (5.22)$$

Notice that the jumps on vertical contours close to the origin, $\mathbf{D}_{\pm}^0(z)$, are exponentially close to the identity matrix.

6. FIRST TRANSFORMATION OF THE RHP

Define the matrix function $\mathbf{T}_n(z)$ as follows from the equation,

$$\mathbf{R}_n(z) = e^{\frac{n}{2}\sigma_3} \mathbf{T}_n(z) e^{n(g(z) - \frac{1}{2})\sigma_3}, \quad (6.1)$$

where the Lagrange multiplier l and the function $g(z)$ are as described in Section 2 and $\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ is the third Pauli matrix. Then $\mathbf{T}_n(z)$ satisfies the following Riemann-Hilbert Problem:

- (1) $\mathbf{T}_n(z)$ is analytic in $\mathbb{C} \setminus \Sigma_R$.
- (2) $\mathbf{T}_{n+}(z) = \mathbf{T}_{n-}(z)j_T(z)$ for $z \in \Sigma_R$, where

$$j_T(z) = \begin{cases} e^{n(g_-(z)-\frac{l}{2})\sigma_3} j_R(z) e^{-n(g_+(z)-\frac{l}{2})\sigma_3} & \text{for } z \in \mathbb{R} \\ e^{n(g(z)-\frac{l}{2})\sigma_3} j_R(z) e^{-n(g(z)-\frac{l}{2})\sigma_3} & \text{for } z \in \Sigma_R \setminus \mathbb{R}. \end{cases} \quad (6.2)$$

(3)

$$\mathbf{T}_n(z) \sim \mathbf{I} + \frac{\mathbf{T}_1}{z} + \frac{\mathbf{T}_2}{z^2} + \dots, \quad \text{as } z \rightarrow \infty. \quad (6.3)$$

From (2.23) we have that

$$g(z) = \log z + O(z^{-1}), \quad \text{as } z \rightarrow \infty. \quad (6.4)$$

This implies that

$$[\mathbf{T}_1]_{12} = e^{-nl}[\mathbf{R}_1]_{12}. \quad (6.5)$$

Let's take a closer look at the behavior of the jump matrix j_T described in (6.2) on the horizontal segments of Σ_R . We have that

$$j_T(z) = \begin{cases} \begin{pmatrix} e^{-nG(z)} & e^{n(g_+(z)+g_-(z)-V(z)-l)} \\ 0 & e^{nG(z)} \end{pmatrix} & \text{when } z \in (-\infty, \alpha') \cup (\beta', \infty), \\ \begin{pmatrix} e^{-nG(z)} & 0 \\ -(\frac{n\pi}{\gamma})^2 e^{-n(g_+(z)+g_-(z)-V(z)-l)} & e^{nG(z)} \end{pmatrix} & \text{when } z \in (\alpha', \beta'), \\ \begin{pmatrix} 1 & \pm \frac{e^{\pm nG(z)}}{1-e^{\frac{\mp i n \pi x}{\gamma}} e^{\frac{\varepsilon n \pi x}{\gamma}}} \\ 0 & 1 \end{pmatrix} & \text{when } z = x \pm i\varepsilon \in (\alpha, \alpha') \cup (\beta', \beta) \pm i\varepsilon, \\ \begin{pmatrix} 1 & \pm \frac{e^{n(2g(z)-l-V(z))}}{1-e^{\frac{\mp i n \pi x}{\gamma}} e^{\frac{\varepsilon n \pi x}{\gamma}}} \\ 0 & 1 \end{pmatrix} & \text{when } z = x \pm i\varepsilon \in (-\infty, \alpha) \cup (\beta, \infty) \pm i\varepsilon, \\ \begin{pmatrix} \Pi(z)^{-1} & 0 \\ \frac{i n \pi}{\gamma} e^{\pm \frac{i n \pi x}{2\gamma}} e^{-n(2g(z)-V(z)-l)} & \Pi(z) \end{pmatrix} & \text{when } z \in (\alpha', \beta') \pm i\varepsilon. \end{cases} \quad (6.6)$$

According to the properties of the g -function, we have the following proposition:

Proposition 6.1. *The jump function j_T has the following large n asymptotics:*

$$j_T(z) = \begin{cases} \begin{pmatrix} e^{-nG(z)} & 0 \\ O(e^{-nC(z)}) & e^{nG(z)} \end{pmatrix} & \text{for } z \in (\alpha', \beta'), \\ \begin{pmatrix} e^{-nG(z)} & 1 \\ 0 & e^{nG(z)} \end{pmatrix} & \text{for } z \in (\alpha, \alpha') \cup (\beta', \beta), \\ \begin{pmatrix} 1 & O(e^{-nC(z)}) \\ 0 & 1 \end{pmatrix} & \text{for } z \in (-\infty, \alpha) \cup (\beta, \infty), \\ \begin{pmatrix} 1 & e^{\pm nG(z)} O(e^{-\frac{\varepsilon n \pi}{\gamma}}) \\ 0 & 1 \end{pmatrix} & \text{for } z \in (\alpha, \alpha') \cup (\beta', \beta) \pm i\varepsilon, \end{cases} \quad (6.7)$$

where $C(z)$ is a positive continuous function on any subset of the given interval which is bounded away from the endpoints of each interval and satisfies

$$C(z) > c|z + 1| \text{ for some } c > 0. \quad (6.8)$$

7. SECOND TRANSFORMATION OF THE RHP

The second transformation is based on two observations. The first is the well known “opening of the lenses” in a neighborhood of the unconstrained support of the equilibrium measure. Namely, notice that, for $x \in (\alpha, \alpha') \cup (\beta', \beta)$, the jump matrix $j_T(x)$ factorizes as

$$\begin{aligned} j_T(x) &= \begin{pmatrix} e^{-nG(x)} & 1 \\ 0 & e^{nG(x)} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ e^{nG(x)} & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ e^{-nG(x)} & 1 \end{pmatrix} \\ &= j_-(x)j_Mj_+(x), \end{aligned} \quad (7.1)$$

which allows us to reduce the jump matrix j_T to the one j_M plus asymptotically small jumps on the lens boundaries. The second observation consists of two facts. Firstly, the jumps on the segments $[\alpha', \beta'] \pm i\varepsilon$ behave, for large n , as $\pm e^{\pm \frac{in\pi z}{2\gamma}}$. Secondly, note that, for $x \in [\alpha', \beta']$, $G(x)$ is a linear function with slope $-\frac{\pi i}{\gamma}$. With these facts in mind, we make the second transformation of the RHP. Let

$$\mathbf{S}_n(z) = \begin{cases} \mathbf{T}_n(z)j_+(z)^{-1} & \text{for } z \in \{(\alpha, \alpha') \cup (\beta', \beta)\} \times (0, i\varepsilon) \\ \mathbf{T}_n(z)j_-(z) & \text{for } z \in \{(\alpha, \alpha') \cup (\beta', \beta)\} \times (0, -i\varepsilon) \\ \mathbf{T}_n(z) \begin{pmatrix} -\frac{\gamma}{n\pi i} e^{-\frac{in\pi z}{2\gamma}} & 0 \\ 0 & -\frac{n\pi i}{\gamma} e^{\frac{in\pi z}{2\gamma}} \end{pmatrix} & \text{for } z \in (\alpha', \beta') \times (0, i\varepsilon) \\ \mathbf{T}_n(z) \begin{pmatrix} \frac{\gamma}{n\pi i} e^{\frac{in\pi z}{2\gamma}} & 0 \\ 0 & \frac{n\pi i}{\gamma} e^{-\frac{in\pi z}{2\gamma}} \end{pmatrix} & \text{for } z \in (\alpha', \beta') \times (0, -i\varepsilon) \\ \mathbf{T}_n(z) & \text{otherwise.} \end{cases} \quad (7.2)$$

This function satisfies a similar RHP to $\mathbf{T}$, but jumps now occur on a new contour, Σ_S , which is obtained from Σ_R by adding the two segments $(\alpha - i\varepsilon, \alpha + i\varepsilon)$ and $(\beta - i\varepsilon, \beta + i\varepsilon)$, see Fig. 9.




FIGURE 9. The contour Σ_S .

On the horizontal segments for which the jump function j_S differs from j_T , we have that, as $n \rightarrow \infty$,

$$j_S(z) = \begin{cases} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} & \text{for } z \in (\alpha, \alpha') \cup (\beta', \beta) \\ \begin{pmatrix} 1 + O(e^{-\varepsilon n \pi / \gamma}) & O(e^{n[G(z) - \varepsilon \pi / \gamma]}) \\ -e^{-nG(z)} & 1 \end{pmatrix} & \text{for } z - i\varepsilon \in (\alpha, \alpha') \cup (\beta', \beta) \\ \begin{pmatrix} 1 + O(e^{-\varepsilon n \pi / \gamma}) & O(e^{n[-G(z) - \varepsilon n \pi / \gamma]}) \\ e^{nG(z)} & 1 \end{pmatrix} & \text{for } z + i\varepsilon \in (\alpha, \alpha') \cup (\beta', \beta) \\ \begin{pmatrix} 1 + O(e^{-\varepsilon n \pi / \gamma}) & 0 \\ \frac{n\pi i}{\gamma} e^{-n(2g(z) - l - V(z))} & 1 + O(e^{-\varepsilon n \pi / \gamma}) \end{pmatrix} & \text{for } z \in [\alpha', \beta'] \pm i\varepsilon \\ \begin{pmatrix} -e^{-n\pi i(1+\zeta)} & 0 \\ e^{-n(g_+(z) + g_-(z) - l - V(z))} & -e^{n\pi i(1+\zeta)} \end{pmatrix} & \text{for } z \in [\alpha', \beta']. \end{cases} \quad (7.3)$$

By formula (2.34) for the G -function and the upper constraint (2.10) on the density ρ , we obtain that, for sufficiently small $\varepsilon > 0$,

$$0 < \mp \operatorname{Re} G(x \pm i\varepsilon) = 2\pi\rho(x) + O(\varepsilon^2) < \frac{\pi\varepsilon}{\gamma} + O(\varepsilon^2). \quad (7.4)$$

This, combined with property (2.32) of the g -function, imply that all jumps on horizontal segments are exponentially close to the identity matrix, provided that they are bounded away from the interval $[\alpha, \beta]$. For what follows we denote

$$\Omega_n = \pi + n2\pi \int_0^\beta \rho(x) dx = \pi + n\pi(1 + \zeta), \quad (7.5)$$

so that

$$-e^{-n\pi i(1+\zeta)} = e^{-i\Omega_n}. \quad (7.6)$$

8. MODEL RHP

The model RHP appears when we drop in the jump matrix $j_S(z)$ the terms that vanish as $n \rightarrow \infty$:

- (1) $\mathbf{M}(z)$ is analytic in $\mathbb{C} \setminus [\alpha, \beta]$.
- (2) $\mathbf{M}_+(z) = \mathbf{M}_-(z)j_M(z)$ for $z \in [\alpha, \beta]$, where

$$j_M(z) = \begin{cases} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} & \text{for } z \in [\alpha, \alpha'] \cup [\beta, \beta'] \\ e^{-i\Omega_n \sigma_3} & \text{for } z \in [\alpha', \beta']. \end{cases} \quad (8.1)$$

(3)

$$\mathbf{M}(z) \sim \mathbf{I} + \frac{\mathbf{M}_1}{z} + \frac{\mathbf{M}_2}{z^2} + \dots \quad \text{as } z \rightarrow \infty. \quad (8.2)$$

This model problem was first solved, in the general multi-cut case, in [12], and is solved in two steps. In the first step, we solve the following auxiliary RHP:

- (1) $\mathbf{Q}(z)$ is analytic in $\mathbb{C} \setminus [\alpha, \alpha'] \cup [\beta', \beta]$,
- (2) $\mathbf{Q}_+(z) = \mathbf{Q}_-(z) \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ for $z \in [\alpha, \alpha'] \cup [\beta', \beta]$

(3) $\mathbf{Q}(z) = \mathbf{I} + O(z^{-1})$ as $z \rightarrow \infty$.

This RHP has the unique solution (see [12])

$$\mathbf{Q}(z) = \begin{pmatrix} \frac{\gamma(z)+\gamma^{-1}(z)}{2} & \frac{\gamma(z)-\gamma^{-1}(z)}{-2i} \\ \frac{\gamma(z)-\gamma^{-1}(z)}{2i} & \frac{\gamma(z)+\gamma^{-1}(z)}{2} \end{pmatrix} \quad (8.3)$$

where

$$\gamma(z) = \left(\frac{(z-\alpha)(z-\beta')}{(z-\alpha')(z-\beta)} \right)^{1/4} \quad (8.4)$$

with cuts on $[\alpha, \alpha'] \cup [\beta', \beta]$.

To solve the model RHP described in (8.1) and (8.2), we again use elliptic functions. Define the function

$$f(s) = \frac{\vartheta_3(s+d+c)}{\vartheta_3(s+d)} \quad (8.5)$$

where ϑ_3 is as defined in (1.38) with elliptic nome $q = e^{i\pi\tau} = e^{\frac{-\pi^2}{2\gamma}}$ ($\tau = \frac{i\pi}{2\gamma}$), and d and c are arbitrary complex numbers. Notice that f has the following periodic properties:

$$f(s+\pi) = f(s); \quad f(s+\pi\tau) = e^{-2ic} f(s) \quad (8.6)$$

and that f is an even function. Now let

$$\tilde{u}(z) = \frac{\pi}{2K} u(z) = \frac{\pi}{2} \int_{\beta}^z \frac{dz'}{\sqrt{R(z')}} \quad (8.7)$$

where u is as defined in (3.1). Then $\tilde{u}$ is two-valued on $[\alpha, \beta]$ and satisfies

$$\tilde{u}_+(x) - \tilde{u}_-(x) = \pi\tau \quad \text{for } x \in [\alpha', \beta']. \quad (8.8)$$

Also,

$$\tilde{u}_{\pm}(\alpha) = \frac{\pi}{2}, \quad \tilde{u}_{\pm}(\alpha') = \frac{\pi}{2} \pm \frac{\pi\tau}{2}, \quad \tilde{u}_{\pm}(\beta') = \pm \frac{\pi\tau}{2}, \quad \tilde{u}_{\pm}(\beta) = 0, \quad (8.9)$$

cf. Fig. 6. Because $\sqrt{R(x)}_+ = -\sqrt{R(x)}_-$ for $x \in [\alpha, \alpha'] \cup [\beta', \beta]$, it immediately follows that

$$\tilde{u}_+(x) + \tilde{u}_-(x) = 0 \quad \text{for } x \in [\beta', \beta], \quad (8.10)$$

and that

$$\tilde{u}_+(x) + \tilde{u}_-(x) = \tilde{u}_+(\alpha') - \tilde{u}_+(\beta') + \tilde{u}_-(\alpha') - \tilde{u}_-(\beta') = \pi \quad \text{for } x \in [\alpha, \alpha']. \quad (8.11)$$

We now define

$$f_1(z) = \frac{\vartheta_3(\tilde{u}(z) + d + \frac{\Omega_n}{2})}{\vartheta_3(\tilde{u}(z) + d)}, \quad f_2(z) = \frac{\vartheta_3(-\tilde{u}(z) + d + \frac{\Omega_n}{2})}{\vartheta_3(-\tilde{u}(z) + d)} \quad \text{for } z \in \mathbb{C} \setminus [\alpha, \beta] \quad (8.12)$$

where d is an arbitrary complex number. It then follows from (8.6) and (8.8) that

$$f_{1+}(x) = e^{-i\Omega_n} f_{1-}(x) \quad \text{and} \quad f_{2+}(x) = e^{i\Omega_n} f_{2-}(x) \quad \text{for } x \in [\alpha', \beta'], \quad (8.13)$$

and from (8.6), (8.10), and (8.11) that

$$f_{1+}(x) = f_{2-}(x) \quad \text{and} \quad f_{2+}(x) = f_{1-}(x) \quad \text{for } x \in [\alpha, \alpha'] \cup [\beta', \beta]. \quad (8.14)$$

Define the matrix valued function

$$\mathbf{F}(z) = \begin{pmatrix} \frac{\vartheta_3(\tilde{u}(z)+d_1+\frac{\Omega_n}{2})}{\vartheta_3(\tilde{u}(z)+d_1)} & \frac{\vartheta_3(-\tilde{u}(z)+d_1+\frac{\Omega_n}{2})}{\vartheta_3(-\tilde{u}(z)+d_1)} \\ \frac{\vartheta_3(\tilde{u}(z)+d_2+\frac{\Omega_n}{2})}{\vartheta_3(\tilde{u}(z)+d_2)} & \frac{\vartheta_3(-\tilde{u}(z)+d_2+\frac{\Omega_n}{2})}{\vartheta_3(-\tilde{u}(z)+d_2)} \end{pmatrix} \quad (8.15)$$

where d_1 and d_2 are yet undetermined complex constants. Then, from (8.13) and (8.14) we have that

$$\begin{aligned} \mathbf{F}_+(x) &= \mathbf{F}_-(x) \begin{pmatrix} e^{-i\Omega_n} & 0 \\ 0 & e^{i\Omega_n} \end{pmatrix} \quad \text{for } x \in [\alpha', \beta'] \\ \mathbf{F}_+(x) &= \mathbf{F}_-(x) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{for } x \in [\alpha, \alpha'] \cup [\beta', \beta]. \end{aligned} \quad (8.16)$$

We can now combine (8.3) and (8.15) to obtain

$$\mathbf{M}(z) = \mathbf{F}(\infty)^{-1} \begin{pmatrix} \frac{\gamma(z)+\gamma^{-1}(z)}{2} \frac{\vartheta_3(\tilde{u}(z)+d_1+\frac{\Omega_n}{2})}{\vartheta_3(\tilde{u}(z)+d_1)} & \frac{\gamma(z)-\gamma^{-1}(z)}{-2i} \frac{\vartheta_3(-\tilde{u}(z)+d_1+\frac{\Omega_n}{2})}{\vartheta_3(-\tilde{u}(z)+d_1)} \\ \frac{\gamma(z)-\gamma^{-1}(z)}{2i} \frac{\vartheta_3(\tilde{u}(z)+d_2+\frac{\Omega_n}{2})}{\vartheta_3(\tilde{u}(z)+d_2)} & \frac{\gamma(z)+\gamma^{-1}(z)}{2} \frac{\vartheta_3(-\tilde{u}(z)+d_2+\frac{\Omega_n}{2})}{\vartheta_3(-\tilde{u}(z)+d_2)} \end{pmatrix} \quad (8.17)$$

where

$$\mathbf{F}(\infty) = \begin{pmatrix} \frac{\vartheta_3(\tilde{u}_\infty+d_1+\frac{\Omega_n}{2})}{\vartheta_3(\tilde{u}_\infty+d_1)} & 0 \\ 0 & \frac{\vartheta_3(-\tilde{u}_\infty+d_2+\frac{\Omega_n}{2})}{\vartheta_3(-\tilde{u}_\infty+d_2)} \end{pmatrix}. \quad (8.18)$$

and $\tilde{u}_\infty \equiv \tilde{u}(\infty)$. This matrix satisfies conditions (8.1) and (8.2) of the model RHP, but may not be analytic on $\mathbb{C} \setminus [\alpha, \beta]$, as it may have some poles at the zeroes of $\vartheta_3(\pm\tilde{u}(z) + d_{1,2})$. However, we can choose the constants d_1 and d_2 such that these zeroes coincide with the zeroes of $\gamma(z) \pm \gamma^{-1}(z)$ and are thus cancelled in the product.

First consider the zeroes of $\gamma(z) \pm \gamma^{-1}(z)$. These are the zeroes of $\gamma^2(z) \pm 1$ and thus of $\gamma^4(z) - 1$, thus there is only one zero, which uniquely solves the equation

$$p(z) \equiv \frac{(z-\alpha)(z-\beta')}{(z-\alpha')(z-\beta)} = 1, \quad (8.19)$$

which is

$$x_0 = \frac{\beta\alpha' - \alpha\beta'}{(\alpha' - \alpha) + (\beta - \beta')} \in (\alpha', \beta'). \quad (8.20)$$

It is easy to check that $\gamma(x_0) = 1$, thus x_0 is the unique zero of $\gamma(z) - \gamma^{-1}(z)$, whereas there are no zeroes of $\gamma(z) + \gamma^{-1}(z)$ on the specified sheet. We use here the change of variables v defined in (3.2). Notice that, by (3.19),

$$v(x_0) = \frac{\beta' - \alpha}{\beta' - \alpha'} = \frac{\text{dn}^2(u_\infty)}{k^2 \text{cn}^2(u_\infty)} \quad (8.21)$$

implying that

$$\text{sn}^2(u(x_0)) = \frac{\text{dn}^2(u_\infty)}{k^2 \text{cn}^2(u_\infty)}. \quad (8.22)$$

Since $x_0 \in (\alpha', \beta')$, we must have $u(x_0) \in (iK', K + iK')$ (if we choose to take u_+). Since sn^2 is a one-to-one function on this interval there is a unique point $u_0 \in (iK', K + iK')$ such

that $\text{sn}^2(u_0) = \frac{\text{dn}^2(u_\infty)}{k^2 \text{cn}^2(u_\infty)}$. The simple period identity

$$\text{sn}(u + K + iK') = \frac{\text{dn}(u)}{k \text{cn}(u)} \quad (8.23)$$

along with (8.22) gives that we must have

$$u_0 = u(x_0) = K - u_\infty + iK' \quad (8.24)$$

thus

$$\tilde{u}(x_0) = \frac{\pi}{2K}(K - u_\infty + iK') = \frac{\tau\pi}{2} + \frac{\pi}{2} - \tilde{u}_\infty. \quad (8.25)$$

We now consider zeroes of the function $\vartheta_3(\tilde{u}(z) - d) \equiv \vartheta_3(-\tilde{u}(z) + d)$. The zeroes of this function are the solutions to the equation

$$\tilde{u}(z) - d = (2m + 1)\frac{\pi}{2} + (2k + 1)\frac{\tau\pi}{2} \quad (8.26)$$

for any $m, k \in \mathbb{Z}$. Because $\tilde{u}$ maps the first sheet of X to the rectangular domain $[0, \frac{\pi}{2}] \times [-\frac{\tau\pi}{2}, \frac{\tau\pi}{2}]$, it is clear that this equation can have at most one solution, and without any loss of generality we may take $m = k = 0$. Then, if we want the solution of this equation to be x_0 , we need to let

$$d = \tilde{u}(x_0) - \frac{\pi}{2}(1 + \tau) = -\tilde{u}_\infty. \quad (8.27)$$

This choice of d also ensures that $\vartheta_3(\tilde{u}(z) + d) \equiv \vartheta_3(-\tilde{u}(z) - d)$ has no zeroes on the first sheet of X . We can then let

$$d_1 = d, \quad d_2 = -d \quad (8.28)$$

so that (8.17) and (8.18) become

$$\begin{aligned} \mathbf{M}(z) &= \mathbf{F}(\infty)^{-1} \begin{pmatrix} \frac{\gamma(z) + \gamma^{-1}(z)}{2} \frac{\vartheta_3(\tilde{u}(z) + d + \frac{\Omega n}{2})}{\vartheta_3(\tilde{u}(z) + d)} & \frac{\gamma(z) - \gamma^{-1}(z)}{-2i} \frac{\vartheta_3(-\tilde{u}(z) + d + \frac{\Omega n}{2})}{\vartheta_3(-\tilde{u}(z) + d)} \\ \frac{\gamma(z) - \gamma^{-1}(z)}{2i} \frac{\vartheta_3(\tilde{u}(z) - d + \frac{\Omega n}{2})}{\vartheta_3(\tilde{u}(z) - d)} & \frac{\gamma(z) + \gamma^{-1}(z)}{2} \frac{\vartheta_3(-\tilde{u}(z) - d + \frac{\Omega n}{2})}{\vartheta_3(-\tilde{u}(z) - d)} \end{pmatrix} \\ &= \mathbf{F}(\infty)^{-1} \begin{pmatrix} \frac{\gamma(z) + \gamma^{-1}(z)}{2} \frac{\vartheta_3(\tilde{u}(z) + (n + \frac{1}{2})\omega)}{\vartheta_4(\tilde{u}(z) + \frac{\omega}{2})} & \frac{\gamma(z) - \gamma^{-1}(z)}{-2i} \frac{\vartheta_3(\tilde{u}(z) - (n + \frac{1}{2})\omega)}{\vartheta_4(\tilde{u}(z) - \frac{\omega}{2})} \\ \frac{\gamma(z) - \gamma^{-1}(z)}{2i} \frac{\vartheta_3(\tilde{u}(z) + (n - \frac{1}{2})\omega)}{\vartheta_4(\tilde{u}(z) - \frac{\omega}{2})} & \frac{\gamma(z) + \gamma^{-1}(z)}{2} \frac{\vartheta_3(\tilde{u}(z) - (n - \frac{1}{2})\omega)}{\vartheta_4(\tilde{u}(z) + \frac{\omega}{2})} \end{pmatrix} \end{aligned} \quad (8.29)$$

where

$$\mathbf{F}(\infty) = \begin{pmatrix} \frac{\vartheta_3(\frac{\Omega n}{2})}{\vartheta_3(0)} & 0 \\ 0 & \frac{\vartheta_3(\frac{\Omega n}{2})}{\vartheta_3(0)} \end{pmatrix} = \begin{pmatrix} \frac{\vartheta_4(n\omega)}{\vartheta_3(0)} & 0 \\ 0 & \frac{\vartheta_4(n\omega)}{\vartheta_3(0)} \end{pmatrix}, \quad (8.30)$$

solving the model RHP. The asymptotics at infinity are

$$\mathbf{M}(z) = I + \frac{\mathbf{M}_1}{z} + O(z^{-2}) \quad (8.31)$$

where the matrix $\mathbf{M}_1$ has the form

$$\mathbf{M}_1 = \begin{pmatrix} * & \frac{\vartheta_3(-\tilde{u}_\infty + d + \frac{\Omega n}{2})\vartheta_3(\tilde{u}_\infty + d)}{\vartheta_3(\tilde{u}_\infty + d + \frac{\Omega n}{2})\vartheta_3(-\tilde{u}_\infty + d)} \frac{(\beta - \beta') + (\alpha' - \alpha)}{-4i} \\ \frac{\vartheta_3(\tilde{u}_\infty - d + \frac{\Omega n}{2})\vartheta_3(-\tilde{u}_\infty - d)}{\vartheta_3(-\tilde{u}_\infty - d + \frac{\Omega n}{2})\vartheta_3(\tilde{u}_\infty - d)} \frac{(\beta - \beta') + (\alpha' - \alpha)}{4i} & * \end{pmatrix}. \quad (8.32)$$

The matrix M_1 can be written in a cleaner fashion and in terms of the original parameters as follows:

Proposition 8.1. *We have that*

$$[\mathbf{M}_1]_{12} = \frac{iA(\omega)\vartheta_4((n+1)\omega)}{\vartheta_4(n\omega)}, \quad [\mathbf{M}_1]_{21} = \frac{A(\omega)\vartheta_4(n\omega)}{i\vartheta_4((n-1)\omega)}. \quad (8.33)$$

where

$$\omega = \frac{\pi(1+\zeta)}{2}, \quad A(\omega) = \frac{\pi\vartheta_1'(0)}{2\vartheta_1(\omega)}. \quad (8.34)$$

For a proof of this proposition see Appendix D in the end of the paper. Notice that since $\mathbf{M}$ solves the model RHP, we have that

$$\det \mathbf{M}(z) = 1, \quad z \in \mathbb{C}. \quad (8.35)$$

9. PARAMETRIX AT OUTER TURNING POINTS

We now consider small disks $D(\alpha, \varepsilon)$ and $D(\beta, \varepsilon)$ centered at the outer turning points. Denote $D = D(\alpha, \varepsilon) \cup D(\beta, \varepsilon)$. We will seek a local parametrix $\mathbf{U}_n(z)$ defined on D such that

$$(1) \quad \mathbf{U}_n(z) \text{ is analytic on } D \setminus \Sigma_S. \quad (9.1)$$

$$(2) \quad \mathbf{U}_{n+}(z) = \mathbf{U}_{n-}(z)j_S(z) \quad \text{for } z \in D \cap \Sigma_S. \quad (9.2)$$

$$(3) \quad \mathbf{U}_n(z) = \mathbf{M}(z)(I + O(n^{-1})) \quad \text{uniformly for } z \in \partial D. \quad (9.3)$$

We first construct the parametrix near β . The jumps j_S are given by

$$j_S(z) = \begin{cases} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} & \text{for } z \in (\beta - \varepsilon, \beta), \\ \begin{pmatrix} 1 & 0 \\ -e^{-nG(z)} & 1 \end{pmatrix} & \text{for } z \in (\beta, \beta + i\varepsilon), \\ \begin{pmatrix} 1 & 0 \\ e^{nG(z)} & 1 \end{pmatrix} & \text{for } z \in (\beta, \beta - i\varepsilon), \\ \begin{pmatrix} e^{-nG(z)} & e^{n(g_+(z)+g_-(z)-V(z)-l)} \\ 0 & e^{nG(z)} \end{pmatrix} & \text{for } z \in (\beta, \beta + \varepsilon). \end{cases} \quad (9.4)$$

If we let

$$\mathbf{U}_n(z) = \mathbf{Q}_n(z)e^{-n(g(z)-\frac{V(z)}{2}-\frac{l}{2})\sigma_3}, \quad (9.5)$$

then the jump conditions on $\mathbf{Q}_n$ become

$$\mathbf{Q}_{n+}(z) = \mathbf{Q}_{n-}(z)j_Q(z) \quad (9.6)$$

where

$$j_Q(z) = \begin{cases} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} & \text{for } z \in (\beta - \varepsilon, \beta), \\ \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix} & \text{for } z \in (\beta, \beta + i\varepsilon), \\ \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} & \text{for } z \in (\beta, \beta - i\varepsilon), \\ \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} & \text{for } z \in (\beta, \beta + \varepsilon). \end{cases} \quad (9.7)$$

where orientation is from left to right on horizontal contours, and down to up on vertical contours, according to Figure 7.

$\mathbf{Q}_n$ can be constructed using Airy functions. The Airy function solves the differential equation $y'' = zy$, and has the following asymptotics at infinity:

$$\begin{aligned} \text{Ai}(z) &= \frac{1}{2\sqrt{\pi}z^{1/4}} e^{-\frac{2}{3}z^{3/2}} \left(1 - \frac{5}{48}z^{-3/2} + O(z^{-3}) \right) \\ \text{Ai}'(z) &= -\frac{1}{2\sqrt{\pi}} z^{1/4} e^{-\frac{2}{3}z^{3/2}} \left(1 + \frac{7}{48}z^{-3/2} + O(z^{-3}) \right) \end{aligned} \quad (9.8)$$

as $z \rightarrow \infty$ with $\arg z \in (-\pi + \varepsilon, \pi - \varepsilon)$ for any $\varepsilon > 0$. If we let

$$y_0(z) = \text{Ai}(z), \quad y_1(z) = \omega \text{Ai}(\omega z), \quad y_2(z) = \omega^2 \text{Ai}(\omega^2 z) \quad (9.9)$$

where $\omega = e^{\frac{2\pi i}{3}}$, then the functions y_0, y_1 , and y_2 satisfy the relation

$$y_0(z) + y_1(z) + y_2(z) = 0. \quad (9.10)$$

If we take

$$\Phi_\beta(z) = \begin{cases} \begin{pmatrix} y_0(z) & -y_2(z) \\ y_0'(z) & -y_2'(z) \end{pmatrix} & \text{for } \arg z \in \left(0, \frac{\pi}{2}\right), \\ \begin{pmatrix} -y_1(z) & -y_2(z) \\ -y_1'(z) & -y_2'(z) \end{pmatrix} & \text{for } \arg z \in \left(\frac{\pi}{2}, \pi\right), \\ \begin{pmatrix} -y_2(z) & y_1(z) \\ -y_2'(z) & y_1'(z) \end{pmatrix} & \text{for } \arg z \in \left(-\pi, -\frac{\pi}{2}\right), \\ \begin{pmatrix} y_0(z) & y_1(z) \\ y_0'(z) & y_1'(z) \end{pmatrix} & \text{for } \arg z \in \left(-\frac{\pi}{2}, 0\right), \end{cases} \quad (9.11)$$

then Φ_β satisfies jump conditions similar to (9.7), but for jumps on rays emanating from the origin rather than from β . We thus need to map the disk $D(\beta, \varepsilon)$ onto some convex neighborhood of the origin in order to take advantage of the function Φ_β . Our mapping should match the asymptotics of the Airy function in order to have the matching property (9.3).

To this end notice that, by (2.19), for $t \in [\beta', \beta]$, as $t \rightarrow \beta$,

$$\rho(t) = C(\beta - t)^{1/2} + O((\beta - t)^{3/2}), \quad C > 0. \quad (9.12)$$

It follows that, as $z \rightarrow \beta$ for $z \in (\beta', \beta)$,

$$\int_z^\beta \rho(t) dt = C_0(\beta - z)^{3/2} + O((\beta - z)^{5/2}), \quad C_0 = \frac{2}{3}C. \quad (9.13)$$

Thus

$$\psi_\beta(z) = - \left\{ \frac{3\pi}{2} \int_z^\beta \rho(t) dt \right\}^{2/3} \quad (9.14)$$

is analytic at β , and so extends to a conformal map from $D(\beta, \varepsilon)$ (for small enough ε) onto a convex neighborhood of the origin. Furthermore,

$$\psi_\beta(\beta) = 0, \quad \psi'_\beta(\beta) > 0; \quad (9.15)$$

therefore ψ_β is real negative on $(\beta - \varepsilon, \beta)$, and real positive on $(\beta, \beta + \varepsilon)$. Also, we can slightly deform the vertical pieces of the contour Σ_S close to β , so that

$$\psi_\beta\{D(\beta, \varepsilon) \cap \Sigma_S\} = (-\varepsilon, \varepsilon) \cup (-i\varepsilon, i\varepsilon). \quad (9.16)$$

We now set

$$\mathbf{Q}_n(z) = \mathbf{E}_n^\beta(z) \Phi_\beta(n^{2/3} \psi_\beta(z)) \quad (9.17)$$

so that

$$\mathbf{U}_n(z) = \mathbf{E}_n^\beta(z) \Phi_\beta(n^{2/3} \psi_\beta(z)) e^{-n(g(z) - \frac{V(z)}{2} - \frac{1}{2})\sigma_3} \quad (9.18)$$

where

$$\mathbf{E}_n^\beta(z) = \mathbf{M}(z) \mathbf{L}_n^\beta(z)^{-1}, \quad \mathbf{L}_n^\beta(z) = \frac{1}{2\sqrt{\pi}} \begin{pmatrix} n^{-1/6} \psi_\beta^{-1/4}(z) & 0 \\ 0 & n^{1/6} \psi_\beta^{1/4}(z) \end{pmatrix} \begin{pmatrix} 1 & i \\ -1 & i \end{pmatrix}, \quad (9.19)$$

where we take the branch of $\psi_\beta^{1/4}$ which is positive on $(\beta, \beta + \varepsilon)$ and has a cut on $(\beta - \varepsilon, \beta)$. We claim that $\mathbf{E}_n^\beta(z)$ is analytic in $D(\beta, \varepsilon)$, thus $\mathbf{U}(z)$ has the jump conditions of j_S . This is clear, as both $\mathbf{M}$ and $\mathbf{L}_n^\beta$ have jump the same constant jump, $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$, on the interval $(\beta - \varepsilon, \beta]$, and are analytic elsewhere. The only other possible singularity for either $\mathbf{M}$ or $\mathbf{L}_n^\beta$ is the isolated singularity at β , and this is at most a fourth-root singularity, thus removable. It follows that $\mathbf{E}_n^\beta(z) = \mathbf{M}(z) \mathbf{L}_n^\beta(z)^{-1}$ is analytic on $D(\beta, \varepsilon)$, thus $\mathbf{U}_n$ has the prescribed jumps in $D(\beta, \varepsilon)$.

We are left only to prove the matching condition (9.3). Using (9.8), one can check that, for z in each of the sectors of analyticity, $\Phi_\beta(n^{2/3} \psi_\beta(z))$ satisfies the following asymptotics as $n \rightarrow \infty$:

$$\begin{aligned} \Phi_\beta(n^{2/3} \psi_\beta(z)) &= \frac{1}{2\sqrt{\pi}} n^{\frac{1}{6}\sigma_3} \psi_\beta(z)^{-\frac{1}{4}\sigma_3} \left[\begin{pmatrix} 1 & i \\ -1 & i \end{pmatrix} + \frac{\psi_\beta(z)^{-3/2}}{48n} \begin{pmatrix} -5 & 5i \\ -7 & -7i \end{pmatrix} + O(n^{-2}) \right] \\ &\quad \times e^{-\frac{2}{3}n\psi_\beta(z)^{3/2}\sigma_3} \end{aligned} \quad (9.20)$$

where we always take the principal branch of $\psi_\beta(z)^{3/2}$. As such, $\psi_\beta(z)^{3/2}$ is two-valued for $z \in (\beta - \varepsilon, \beta)$, so that

$$\left[\frac{2}{3} \psi_\beta(x)^{3/2} \right]_\pm = \mp \pi i \int_x^\beta \rho(t) dt. \quad (9.21)$$

Notice that, by (2.32) and (2.35), for $x \in (\beta - \varepsilon, \beta)$,

$$2g_{\pm}(x) - V(x) = l \pm 2\pi i \int_x^{\beta} \rho(t) dt. \quad (9.22)$$

This implies that

$$[2g_{\pm}(\beta) - V(\beta)] - [2g_{\pm}(x) - V(x)] = \mp 2\pi i \int_x^{\beta} \rho(t) dt. \quad (9.23)$$

Combining these equations with (9.21) gives

$$\left[\frac{2}{3} \psi_{\beta}(x)^{3/2} \right]_{\pm} = \frac{1}{2} \left[(2g_{\pm}(\beta) - V(\beta)) - (2g_{\pm}(x) - V(x)) \right]. \quad (9.24)$$

This equation can be extended into the upper and lower planes, respectively, giving

$$\frac{2}{3} \psi_{\beta}(z)^{3/2} = \frac{1}{2} \left[(2g_{\pm}(\beta) - V(\beta)) - (2g(z) - V(z)) \right] \quad \text{for } \pm \operatorname{Im} z > 0. \quad (9.25)$$

Since, by (9.22), $2g_{\pm}(\beta) - V(\beta) = l$, we get that

$$\frac{2}{3} \psi_{\beta}(z)^{3/2} = -g(z) + \frac{V(z)}{2} + \frac{l}{2} \quad (9.26)$$

for z throughout $D(\beta, \varepsilon)$. Plugging (9.20) and (9.26) into (9.18), we get

$$\begin{aligned} \mathbf{U}_n(z) &= \mathbf{M}(z) \mathbf{L}_n^{\beta}(z)^{-1} \frac{1}{2\sqrt{\pi}} n^{-\frac{1}{6}\sigma_3} \psi_{\beta}(z)^{-\frac{1}{4}\sigma_3} \left[\begin{pmatrix} 1 & i \\ -1 & i \end{pmatrix} + \frac{\psi_{\beta}(z)^{-3/2}}{48n} \begin{pmatrix} -5 & 5i \\ -7 & -7i \end{pmatrix} \right. \\ &\quad \left. + O(n^{-2}) \right] e^{n(g(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} e^{-n(g(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} \\ &= \mathbf{M}(z) \left[I + \frac{\psi_{\beta}(z)^{-3/2}}{48n} \begin{pmatrix} 1 & 6i \\ 6i & -1 \end{pmatrix} + O(n^{-2}) \right]. \end{aligned} \quad (9.27)$$

Thus we have that $\mathbf{U}_n$ satisfies conditions (9.1), (9.2), and (9.3).

A similar construction gives the parametrix at the α . Namely, if we let

$$\psi_{\alpha}(z) = - \left\{ \frac{3\pi}{2} \int_{\alpha}^z \rho(t) dt \right\}^{2/3}, \quad (9.28)$$

then ψ_{α} is analytic throughout $D(\alpha, \varepsilon)$, real valued on the real line, and has negative derivative at α . Close to α , the jumps j_Q become

$$j_Q(z) = \begin{cases} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} & \text{for } z \in (\alpha - \varepsilon, \alpha), \\ \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix} & \text{for } z \in (\alpha, \alpha + i\varepsilon), \\ \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} & \text{for } z \in (\alpha, \alpha - i\varepsilon), \\ \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} & \text{for } z \in (\alpha, \alpha + \varepsilon), \end{cases} \quad (9.29)$$

where orientation is taken left to right on horizontal contours, and up to down on vertical contours according to Figure 7. After the change of variables ψ_α (and a slight deformation of vertical contours), these jumps become the following jumps close to the origin:

$$j_Q(\psi_\alpha(z)) = \begin{cases} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} & \text{for } \psi_\alpha(z) \in (-\varepsilon, 0), \\ \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} & \text{for } \psi_\alpha(z) \in (0, i\varepsilon), \\ \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix} & \text{for } \psi_\alpha(z) \in (0, -i\varepsilon), \\ \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} & \text{for } \psi_\alpha(z) \in (0, \varepsilon), \end{cases} \quad (9.30)$$

where orientation is taken right to left on horizontal contours, and down to up on vertical contours. These jump conditions are satisfied by the function

$$\Phi_\alpha(z) = \Phi_\beta(z) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (9.31)$$

Then we can take

$$\mathbf{U}_n(z) = \mathbf{E}_n^\alpha(z) \Phi_\alpha(n^{2/3}\psi_\alpha(z)) e^{-n(g(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} \quad (9.32)$$

for $z \in D(\alpha, \varepsilon)$, where

$$\begin{aligned} \mathbf{E}_n^\alpha(z) &= \mathbf{M}(z) \mathbf{L}_n^\alpha(z)^{-1}, \\ \mathbf{L}_n^\alpha(z) &= \frac{1}{2\sqrt{\pi}} \begin{pmatrix} n^{-1/6}\psi_\alpha^{-1/4}(z) & 0 \\ 0 & n^{1/6}\psi_\alpha^{1/4}(z) \end{pmatrix} \begin{pmatrix} -1 & i \\ 1 & i \end{pmatrix}. \end{aligned} \quad (9.33)$$

Similar to (9.20), we have that in each sector of analyticity, $\Phi_\alpha(n^{2/3}\psi_\alpha(z))$ satisfies

$$\begin{aligned} \Phi_\alpha(n^{2/3}\psi_\alpha(z)) &= \frac{1}{2\sqrt{\pi}} n^{-\frac{1}{6}\sigma_3} \psi_\alpha(z)^{-\frac{1}{4}\sigma_3} \left[\begin{pmatrix} 1 & -i \\ -1 & -i \end{pmatrix} + \frac{\psi_\alpha(z)^{-3/2}}{48n} \begin{pmatrix} -5 & -5i \\ -7 & 7i \end{pmatrix} + O(n^{-2}) \right] \\ &\quad \times e^{-\frac{2}{3}n\psi_\alpha(z)^{3/2}\sigma_3}. \end{aligned} \quad (9.34)$$

Once again, we have that, for $x \in (\alpha, \alpha + \varepsilon)$, $\psi_\alpha(x)^{3/2}$ takes limiting values from above and below, so that

$$\left[\frac{2}{3} \psi_\alpha(x)^{3/2} \right]_{\pm} = \pm \pi i \int_{\alpha}^x \rho(t) dt. \quad (9.35)$$

In analogue to (9.24), we have

$$\frac{2}{3} \psi_\alpha(z)^{3/2} = \frac{1}{2} \left[(2g_{\pm}(\alpha) - V(\alpha)) - (2g(z) - V(z)) \right] \quad \text{for } \pm \operatorname{Im} z > 0. \quad (9.36)$$

Since, by (9.22), $2g_{\pm}(\alpha) - V(\alpha) = l \pm \pi i$, we get that

$$\frac{2}{3} \psi_\alpha(z)^{3/2} = -g(z) + \frac{V(z)}{2} + \frac{l}{2} \pm \pi i \quad \text{for } \pm \operatorname{Im} z > 0. \quad (9.37)$$

Plugging (9.37) into (9.32) and (9.34) gives, as $n \rightarrow \infty$,

$$\begin{aligned}
\mathbf{U}_n(z) &= \mathbf{M}(z) \mathbf{L}_n^\alpha(z)^{-1} \frac{1}{2\sqrt{\pi}} n^{\frac{1}{6}\sigma_3} \psi_\alpha(z)^{-\frac{1}{4}\sigma_3} \left[\begin{pmatrix} 1 & -i \\ -1 & -i \end{pmatrix} + \frac{\psi_\alpha(z)^{-3/2}}{48n} \begin{pmatrix} -5 & -5i \\ -7 & 7i \end{pmatrix} \right. \\
&\quad \left. + O(n^{-2}) \right] e^{n(g(z) - \frac{V(z)}{2} - \frac{1}{2})\sigma_3} \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} e^{-n(g(z) - \frac{V(z)}{2} - \frac{1}{2})\sigma_3} \\
&= \mathbf{M}(z) \begin{pmatrix} -1 & i \\ 1 & i \end{pmatrix}^{-1} \left[\begin{pmatrix} -1 & i \\ 1 & i \end{pmatrix} + \frac{\psi_\alpha(z)^{-3/2}}{48n} \begin{pmatrix} 5 & 5i \\ 7 & -7i \end{pmatrix} + O(n^{-2}) \right] \\
&= \mathbf{M}(z) \left[I + \frac{\psi_\alpha(z)^{-3/2}}{48n} \begin{pmatrix} 1 & -6i \\ -6i & -1 \end{pmatrix} + O(n^{-2}) \right].
\end{aligned} \tag{9.38}$$

10. PARAMETRIX AT THE INNER TURNING POINTS

We now consider small disks $D(\alpha', \varepsilon)$ and $D(\beta', \varepsilon)$ centered at the inner turning points. Denote $\tilde{D} = D(\alpha', \varepsilon) \cup D(\beta', \varepsilon)$. We will seek a local parametrix $\mathbf{U}_n(z)$ defined on $\tilde{D}$ such that

(1)

$$\mathbf{U}_n(z) \text{ is analytic on } \tilde{D} \setminus \Sigma_S. \tag{10.1}$$

(2)

$$\mathbf{U}_{n+}(z) = \mathbf{U}_{n-}(z) j_S(z) \quad \text{for } z \in \tilde{D} \cap \Sigma_S. \tag{10.2}$$

(3)

$$\mathbf{U}_n(z) = \mathbf{M}(z) (I + O(n^{-1})) \quad \text{uniformly for } z \in \partial\tilde{D}. \tag{10.3}$$

We first construct the parametrix near α' . Let

$$\mathbf{U}_n(z) = \tilde{\mathbf{Q}}_n(z) e^{\mp \frac{i n \pi z}{2\gamma} \sigma_3} e^{-n(g(z) - \frac{V(z)}{2} - \frac{1}{2})\sigma_3} \quad \text{for } \pm \operatorname{Im} z > 0. \tag{10.4}$$

Then the jumps for $\tilde{\mathbf{Q}}_n$ are

$$j_{\tilde{Q}}(z) = \begin{cases} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} & \text{for } z \in (\alpha' - \varepsilon, \alpha'), \\ \begin{pmatrix} -1 & 0 \\ -1 & -1 \end{pmatrix} & \text{for } z \in (\alpha', \alpha' + \varepsilon), \\ \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} & \text{for } z \in (\alpha', \alpha' + i\varepsilon), \\ \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} & \text{for } z \in (\alpha', \alpha' - i\varepsilon), \end{cases} \tag{10.5}$$

where orientation is taken from left to right on horizontal contours, and down to up on vertical contours according to Figure 7. A proof of this statement is given in Appendix C.

We now take

$$\Phi_{\alpha'}(z) = \begin{cases} \begin{pmatrix} y_2(z) & -y_0(z) \\ y_2'(z) & -y_0'(z) \end{pmatrix} & \text{for } \arg z \in \left(0, \frac{\pi}{2}\right), \\ \begin{pmatrix} y_2(z) & y_1(z) \\ y_2'(z) & y_1'(z) \end{pmatrix} & \text{for } \arg z \in \left(\frac{\pi}{2}, \pi\right), \\ \begin{pmatrix} y_1(z) & -y_2(z) \\ y_1'(z) & -y_2'(z) \end{pmatrix} & \text{for } \arg z \in \left(-\pi, -\frac{\pi}{2}\right), \\ \begin{pmatrix} y_1(z) & y_0(z) \\ y_1'(z) & y_0'(z) \end{pmatrix} & \text{for } \arg z \in \left(-\frac{\pi}{2}, 0\right). \end{cases} \quad (10.6)$$

Then $\Phi_{\alpha'}(z)$ solves a RHP similar to that of $\tilde{\mathbf{Q}}_n$, but for jumps emanating from the origin rather than from α' .

Notice that, by (2.19), for $t \in [\alpha, \alpha']$, as $t \rightarrow \alpha'$,

$$\rho(t) = \frac{1}{2\gamma} - C(\alpha' - t)^{1/2} + O((\alpha' - t)^{3/2}), \quad C > 0. \quad (10.7)$$

It follows that, as $z \rightarrow \alpha'$, for $z \in (\alpha, \alpha')$,

$$\int_z^{\alpha'} \left(\frac{1}{2\gamma} - \rho(t) \right) dt = C(\alpha' - z)^{3/2} + O((\alpha' - z)^{5/2}), \quad C_0 = \frac{2}{3}C \quad (10.8)$$

Thus,

$$\psi_{\alpha'}(z) = - \left\{ \frac{3\pi}{2} \int_z^{\alpha'} \left(\frac{1}{2\gamma} - \rho(t) \right) dt \right\}^{2/3} \quad (10.9)$$

is analytic at α' , and so extends to a conformal map from $D(\alpha', \varepsilon)$ onto a convex neighborhood of the origin. Furthermore,

$$\psi_{\alpha'}(\alpha') = 0, \quad \psi'_{\alpha'}(\alpha') > 0; \quad (10.10)$$

consequently, $\psi_{\alpha'}$ is real negative on $(\alpha' - \varepsilon, \alpha')$, and real positive on $(\alpha', \alpha' + \varepsilon)$. Again, we can slightly deform the vertical pieces of the contour Σ_S close to α' , so that

$$\psi_{\alpha'}\{D(\alpha', \varepsilon) \cap \Sigma_S\} = (-\varepsilon, \varepsilon) \cup (-i\varepsilon, i\varepsilon) \quad (10.11)$$

We now take

$$\tilde{\mathbf{Q}}_n(z) = \mathbf{E}_n^{\alpha'}(z) \Phi_{\alpha'}(n^{2/3} \psi_{\alpha'}(z)) \quad (10.12)$$

where

$$\begin{aligned} \mathbf{E}_n^{\alpha'}(z) &= \mathbf{M}(z) e^{\pm \frac{i\Omega_n}{2} \sigma_3} \tilde{\mathbf{L}}_n(z)^{-1} \quad \text{for } \pm \operatorname{Im} z \geq 0, \\ \mathbf{L}_n^{\alpha'}(z) &= \frac{1}{2\sqrt{\pi}} \begin{pmatrix} n^{-1/6} \psi_{\alpha'}^{-1/4}(z) & 0 \\ 0 & n^{1/6} \psi_{\alpha'}^{1/4}(z) \end{pmatrix} \begin{pmatrix} 1 & i \\ 1 & -i \end{pmatrix}, \end{aligned} \quad (10.13)$$

and we take the branch of $\psi_{\alpha'}^{1/4}$ which is positive on $(\alpha', \alpha' + \varepsilon)$ and has a cut on $(\alpha' - \varepsilon, \alpha')$. $\mathbf{U}_n$ then becomes

$$\begin{aligned} \mathbf{U}_n(z) &= \mathbf{M}(z) e^{\pm \frac{i\Omega_n}{2} \sigma_3} \mathbf{L}_n^{\alpha'}(z)^{-1} \Phi_{\alpha'}(n^{2/3} \psi_{\alpha'}(z)) e^{\mp \frac{i n \pi z}{2\gamma} \sigma_3} e^{-n(g(z) - \frac{V(z)}{2} - \frac{l}{2}) \sigma_3} \\ &\quad \text{for } \pm \operatorname{Im} z > 0. \end{aligned} \quad (10.14)$$

The function $\Phi_{\alpha'}(n^{2/3}\psi_{\alpha'}(z))$ has the jumps j_S , and we claim that the prefactor $\mathbf{E}_n^{\alpha'}$ is analytic in $D(\alpha', \varepsilon)$, thus does not change these jumps. This can be seen, as

$$\mathbf{M}_+(z)e^{\frac{i\Omega_n}{2}\sigma_3} = \mathbf{M}_-(z)e^{-\frac{i\Omega_n}{2}\sigma_3}e^{\frac{i\Omega_n}{2}\sigma_3}j_Me^{\frac{i\Omega_n}{2}\sigma_3} \quad (10.15)$$

thus the jump for the function $\mathbf{M}(z)e^{\pm\frac{i\Omega_n}{2}\sigma_3}$ is

$$e^{\frac{i\Omega_n}{2}\sigma_3}j_Me^{\frac{i\Omega_n}{2}\sigma_3} = \begin{cases} e^{\frac{i\Omega_n}{2}\sigma_3} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} e^{\frac{i\Omega_n}{2}\sigma_3} & \text{for } z \in (\alpha' - \varepsilon, \alpha'), \\ e^{\frac{i\Omega_n}{2}\sigma_3} e^{-i\Omega_n\sigma_3} e^{\frac{i\Omega_n}{2}\sigma_3} & \text{for } z \in (\alpha', \alpha' + \varepsilon), \end{cases} \quad (10.16)$$

or equivalently,

$$e^{\frac{i\Omega_n}{2}\sigma_3}j_Me^{\frac{i\Omega_n}{2}\sigma_3} = \begin{cases} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} & \text{for } z \in (\alpha' - \varepsilon, \alpha'), \\ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} & \text{for } z \in (\alpha', \alpha' + \varepsilon), \end{cases} \quad (10.17)$$

which is exactly the same as the jump conditions for $\mathbf{L}_n^{\alpha'}$. Thus $\mathbf{E}_n^{\alpha'}(z) = \mathbf{M}(z)e^{\pm\frac{i\Omega_n}{2}\sigma_3}\mathbf{L}_n^{\alpha'}(z)^{-1}$ has no jumps in $D(\alpha', \varepsilon)$. The only other possible singularity for $\mathbf{E}_n^{\alpha'}$ is at α' , and this singularity is at most a fourth root singularity, thus removable. Thus, $\mathbf{E}_n^{\alpha'}$ is analytic in $D(\alpha', \varepsilon)$, and $\tilde{\mathbf{Q}}_n$ has the prescribed jumps.

We left check that $\mathbf{U}_n$ satisfies the matching condition (10.3). The large n asymptotics of $\Phi_{\alpha'}(n^{2/3}\psi_{\alpha'}(z))$ are given in the different regions of analyticity as follows:

$$\begin{aligned} \Phi_{\alpha'}(n^{2/3}\psi_{\alpha'}(z)) &= \frac{1}{2\sqrt{\pi}}n^{-\frac{1}{6}\sigma_3}\psi_{\alpha'}(z)^{-\frac{1}{4}\sigma_3} \left[\mp \begin{pmatrix} i & 1 \\ i & -1 \end{pmatrix} \pm \frac{\psi_{\alpha'}(z)^{-3/2}}{48n} \begin{pmatrix} -5i & 5 \\ 7i & 7 \end{pmatrix} \right. \\ &\quad \left. + O(n^{-2}) \right] e^{\frac{2}{3}n\psi_{\alpha'}(z)^{3/2}\sigma_3} \quad \text{for } \pm \operatorname{Im} z > 0, \end{aligned} \quad (10.18)$$

where we always take the principal branch of $\psi_{\alpha'}(z)^{3/2}$. As such, $\psi_{\alpha'}(z)^{3/2}$ is two-valued for $x \in (\alpha' - \varepsilon, \alpha')$, so that

$$\left[\frac{2}{3}\psi_{\alpha'}(x)^{3/2} \right]_{\pm} = \mp \pi i \int_x^{\alpha'} \left(\frac{1}{2\gamma} - \rho(t) \right) dt = \mp \frac{\pi i}{2\gamma}(\alpha' - x) \pm \pi i \int_x^{\alpha'} \rho(t) dt. \quad (10.19)$$

From (2.32) and (2.35), we have that

$$2g_+(x) - V(x) = l + 2\pi i \int_x^{\beta} \rho(t) dt, \quad 2g_-(x) - V(x) = l - 2\pi i \int_x^{\beta} \rho(t) dt \quad (10.20)$$

for $x \in (\alpha' - \varepsilon, \alpha')$. These equations imply that

$$(2g_{\pm}(x) - V(x)) - (2g_{\pm}(\alpha') - V(\alpha')) = \pm 2\pi i \int_x^{\alpha'} \rho(t) dt. \quad (10.21)$$

We can therefore write (10.19) as

$$\left[\frac{2}{3}\psi_{\alpha'}(x)^{3/2} \right]_{\pm} = \mp \frac{\pi i}{2\gamma}(\alpha' - x) + \frac{1}{2} \left[(2g_{\pm}(x) - V(x)) - (2g_{\pm}(\alpha') - V(\alpha')) \right]. \quad (10.22)$$

We can extend these equations into the upper and lower half-plane, respectively, obtaining

$$\frac{2}{3}\psi_{\alpha'}(z)^{3/2} = \mp \frac{\pi i}{2\gamma}(\alpha' - z) + \frac{1}{2} \left[(2g(z) - V(z)) - (2g_{\pm}(\alpha') - V(\alpha')) \right] \quad \text{for } \pm \operatorname{Im} z > 0. \quad (10.23)$$

Using (10.20) at $x = \alpha'$, we can write

$$\frac{2}{3}\psi_{\alpha'}(z)^{3/2} = \mp \frac{\pi i}{2\gamma}(\alpha' - z) + g(z) - \frac{V(z)}{2} - \frac{l}{2} \mp \pi i \int_{\alpha'}^{\beta} \rho(t) dt \quad \text{for } \pm \operatorname{Im} z > 0, \quad (10.24)$$

or equivalently,

$$\frac{2}{3}\psi_{\alpha'}(z)^{3/2} = g(z) - \frac{V(z)}{2} - \frac{l}{2} \pm \frac{\pi i z}{2\gamma} \mp \frac{i(\Omega_n - \pi)}{2n} \quad \text{for } \pm \operatorname{Im} z > 0. \quad (10.25)$$

Plugging (10.18) and (10.24) into (10.14) gives

$$\begin{aligned} \mathbf{U}_n(z) &= \mathbf{M}(z) e^{\pm \frac{i\Omega_n}{2}\sigma_3} \mathbf{L}_n^{\alpha'}(z)^{-1} \frac{1}{2\sqrt{\pi}} n^{-\frac{1}{6}\sigma_3} \psi_{\alpha'}(z)^{-\frac{1}{4}\sigma_3} \\ &\quad \times \left[\mp \begin{pmatrix} i & 1 \\ i & -1 \end{pmatrix} \pm \frac{\psi_{\alpha'}(z)^{-3/2}}{48n} \begin{pmatrix} -5i & 5 \\ 7i & 7 \end{pmatrix} + O(n^{-2}) \right] \\ &\quad \times e^{n(g(z) - \frac{l}{2} - \frac{V(z)}{2})\sigma_3} e^{\mp \frac{i\Omega_n}{2}\sigma_3} e^{\pm \frac{i\pi}{2}\sigma_3} e^{\pm \frac{in\pi z}{2\gamma}\sigma_3} e^{\mp \frac{in\pi z}{2\gamma}\sigma_3} e^{-n(g(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} \\ &= \mathbf{M}(z) e^{\pm \frac{i\Omega_n}{2}\sigma_3} \mathbf{L}_n^{\alpha'}(z)^{-1} \frac{1}{2\sqrt{\pi}} n^{-\frac{1}{6}\sigma_3} \psi_{\alpha'}(z)^{-\frac{1}{4}\sigma_3} \\ &\quad \times \left[\begin{pmatrix} 1 & i \\ 1 & -i \end{pmatrix} + \frac{\psi_{\alpha'}(z)^{-3/2}}{48n} \begin{pmatrix} 5 & -5i \\ -7 & -7i \end{pmatrix} + O(n^{-2}) \right] e^{\mp \frac{i\Omega_n}{2}\sigma_3} \\ &= \mathbf{M}(z) \left[I + \frac{\psi_{\alpha'}(z)^{-3/2}}{48n} e^{\pm i\frac{\Omega_n}{2}\sigma_3} \begin{pmatrix} -1 & -6i \\ -6i & 1 \end{pmatrix} e^{\mp i\frac{\Omega_n}{2}\sigma_3} + O(n^{-2}) \right] \\ &= \mathbf{M}(z) \left[I + \frac{\psi_{\alpha'}(z)^{-3/2}}{48n} \begin{pmatrix} -1 & -6ie^{\pm i\Omega_n} \\ -6ie^{\mp i\Omega_n} & 1 \end{pmatrix} + O(n^{-2}) \right] \quad \text{for } \pm \operatorname{Im}(z) > 0. \end{aligned} \quad (10.26)$$

We can make a similar construction near β' . Let

$$\psi_{\beta'}(z) = - \left\{ \frac{3\pi}{2} \int_{\beta'}^z \left(\frac{1}{2\gamma} - \rho(t) dt \right) \right\}^{2/3}. \quad (10.27)$$

This function is analytic in $D(\beta', \varepsilon)$ and has negative derivative at β' , thus $\operatorname{Im} z$ and $\operatorname{Im} \psi_{\beta'}(z)$ have opposite signs for $z \in D(\beta', \varepsilon)$. Then the jumps for $\tilde{\mathbf{Q}}_n$ are

$$j_{\tilde{\mathbf{Q}}}(z) = \begin{cases} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} & \text{for } z \in (\beta', \beta' + \varepsilon), \\ \begin{pmatrix} -1 & 0 \\ -1 & -1 \end{pmatrix} & \text{for } z \in (\beta' - \varepsilon, \beta'), \\ \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} & \text{for } z \in (\beta', \beta' + i\varepsilon), \\ \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} & \text{for } z \in (\beta', \beta' - i\varepsilon), \end{cases} \quad (10.28)$$

where the contour is oriented from left to right on horizontal segments and up to down on vertical segments according to Figure 7. After a slight deformation of the vertical contours and the change of variables $\psi_{\beta'}$, these jumps become the following jumps close to the origin:

$$j_{\tilde{Q}}(\psi_{\beta'}(z)) = \begin{cases} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} & \text{for } \psi_{\beta'}(z) \in (-\varepsilon, 0), \\ \begin{pmatrix} -1 & 0 \\ -1 & -1 \end{pmatrix} & \text{for } \psi_{\beta'}(z) \in (0, \varepsilon), \\ \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} & \text{for } \psi_{\beta'}(z) \in (-i\varepsilon, 0), \\ \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} & \text{for } \psi_{\beta'}(z) \in (0, i\varepsilon), \end{cases} \quad (10.29)$$

where the contour is oriented from right to left on horizontal segments and down to up on vertical segments. These jump conditions are satisfied by the function

$$\Phi_{\beta'}(z) = \Phi_{\alpha'}(z) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (10.30)$$

Then we can take for $z \in D(\beta', \varepsilon)$,

$$\mathbf{U}_n(z) = \mathbf{M}(z) e^{\frac{\pm i \Omega_n}{2} \sigma_3} \mathbf{L}_n^{\beta'}(z)^{-1} \Phi_{\beta'}(n^{2/3} \psi_{\beta'}(z)) e^{\mp \frac{i n \pi z}{2\gamma} \sigma_3} e^{-n(g(z) - \frac{V(z)}{2} - \frac{l}{2}) \sigma_3} \quad (10.31)$$

for $\pm \operatorname{Im} z > 0$,

where

$$\mathbf{L}_n^{\beta'}(z) = \frac{1}{2\sqrt{\pi}} \begin{pmatrix} n^{-1/6} \psi_{\beta'}^{-1/4}(z) & 0 \\ 0 & n^{1/6} \psi_{\beta'}^{1/4}(z) \end{pmatrix} \begin{pmatrix} -1 & i \\ -1 & -i \end{pmatrix}. \quad (10.32)$$

We once again have

$$\begin{aligned} \Phi_{\beta'}(n^{2/3} \psi_{\beta'}(z)) &= \frac{1}{2\sqrt{\pi}} n^{-\frac{1}{6} \sigma_3} \psi_{\beta'}(z)^{-\frac{1}{4} \sigma_3} \left[\mp \begin{pmatrix} i & -1 \\ i & 1 \end{pmatrix} \mp \frac{\psi_{\beta'}(z)^{-3/2}}{48n} \begin{pmatrix} 5i & 5 \\ -7i & 7 \end{pmatrix} \right. \\ &\quad \left. + O(n^{-2}) \right] e^{\frac{2}{3} n \psi_{\beta'}(z)^{3/2} \sigma_3} \quad \text{for } \pm \operatorname{Im} \psi_{\beta'}(z) > 0 \text{ (so } \pm \operatorname{Im} z < 0), \end{aligned} \quad (10.33)$$

and for $z \in D(\beta', \varepsilon)$,

$$\frac{2}{3} \psi_{\beta'}^{3/2}(z) = \pm \frac{\pi i z}{2\gamma} + g(z) - \frac{V(z)}{2} - \frac{l}{2} \mp \frac{i(\Omega_n - \pi)}{2n} \quad \text{for } \pm \operatorname{Im} z > 0. \quad (10.34)$$

Combining (10.31), (10.33), and (10.34) gives

$$\begin{aligned}
\mathbf{U}_n(z) &= \mathbf{M}(z) e^{\pm \frac{i\Omega_n}{2}\sigma_3} \mathbf{L}_n^{\beta'}(z)^{-1} \frac{1}{2\sqrt{\pi}} n^{-\frac{1}{6}\sigma_3} \psi_{\beta'}(z)^{-\frac{1}{4}\sigma_3} \\
&\quad \times \left[\pm \begin{pmatrix} i & -1 \\ i & 1 \end{pmatrix} \pm \frac{\psi_{\beta'}(z)^{-3/2}}{48n} \begin{pmatrix} 5i & 5 \\ -7i & 7 \end{pmatrix} + O(n^{-2}) \right] \\
&\quad \times e^{\pm \frac{in\pi z}{2\gamma}\sigma_3} e^{n(g(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} e^{\mp \frac{i\Omega_n}{2}\sigma_3} e^{\pm \frac{i\pi}{2}\sigma_3} e^{\mp \frac{in\pi z}{2\gamma}\sigma_3} e^{-n(g(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} \\
&= \mathbf{M}(z) e^{\pm \frac{i\Omega_n}{2}\sigma_3} \left[I + \frac{\psi_{\beta'}(z)^{-3/2}}{48n} \begin{pmatrix} -1 & 6i \\ 6i & 1 \end{pmatrix} + O(n^{-2}) \right] e^{\mp \frac{i\Omega_n}{2}\sigma_3} \\
&= \mathbf{M}(z) \left[I + \frac{\psi_{\beta'}(z)^{-3/2}}{48n} \begin{pmatrix} -1 & 6ie^{\pm i\Omega_n} \\ 6ie^{\mp i\Omega_n} & 1 \end{pmatrix} + O(n^{-2}) \right] \quad \text{for } \pm \operatorname{Im} z > 0.
\end{aligned} \tag{10.35}$$

11. THE THIRD AND FINAL TRANSFORMATION OF THE RHP

We now consider the contour Σ_X , which consists of the circles $\partial D(\alpha, \varepsilon)$, $\partial D(\alpha', \varepsilon)$, $\partial D(\beta', \varepsilon)$, and $\partial D(\beta, \varepsilon)$, all oriented counterclockwise, together with the parts of $\Sigma_S \setminus ([\alpha, \alpha'] \cup [\beta', \beta])$ which lie outside of the disks $D(\alpha, \varepsilon)$, $D(\alpha', \varepsilon)$, $D(\beta', \varepsilon)$, and $D(\beta, \varepsilon)$, see Fig. 10.




FIGURE 10. The contour Σ_X .

We let

$$\mathbf{X}_n(z) = \begin{cases} \mathbf{S}_n(z) \mathbf{M}(z)^{-1} & \text{for } z \text{ outside the disks } D(\alpha, \varepsilon), D(\alpha', \varepsilon), D(\beta', \varepsilon), D(\beta, \varepsilon), \\ \mathbf{S}_n(z) \mathbf{U}_n(z)^{-1} & \text{for } z \text{ inside the disks } D(\alpha, \varepsilon), D(\alpha', \varepsilon), D(\beta', \varepsilon), D(\beta, \varepsilon). \end{cases} \tag{11.1}$$

Then $\mathbf{X}_n(z)$ solves the following RHP:

- (1) $\mathbf{X}_n(z)$ is analytic on $\mathbb{C} \setminus \Sigma_X$.
- (2) $\mathbf{X}_n(z)$ has the jump properties

$$\mathbf{X}_{n+}(x) = \mathbf{X}_{n-}(x) j_X(x) \tag{11.2}$$

where

$$j_X(z) = \begin{cases} \mathbf{M}(z) \mathbf{U}_n(z)^{-1} & \text{for } z \text{ on the circles} \\ \mathbf{M}(z) j_S \mathbf{M}(z)^{-1} & \text{otherwise.} \end{cases} \tag{11.3}$$

- (3) As $z \rightarrow \infty$,

$$\mathbf{X}_n(z) \sim \mathbf{I} + \frac{\mathbf{X}_1}{z} + \frac{\mathbf{X}_2}{z^2} + \dots \tag{11.4}$$

Additionally, we have that $j_X(z)$ is uniformly close to the identity in the following sense:

$$j_X(z) = \begin{cases} I + O(n^{-1}) & \text{uniformly on the circles} \\ I + O(e^{-C(z)n}) & \text{on the rest of } \Sigma_X, \end{cases} \quad (11.5)$$

where $C(z)$ is a positive, continuous function satisfying (6.8). If we set

$$j_X^0(z) = j_X(z) - I, \quad (11.6)$$

then (11.5) becomes

$$j_X^0(z) = \begin{cases} O(n^{-1}) & \text{uniformly on the circles} \\ O(e^{-C(z)n}) & \text{on the rest of } \Sigma_X. \end{cases} \quad (11.7)$$

The solution to the RHP for $\mathbf{X}_n$ is based on the following lemma:

Lemma 11.1. *Suppose $v(z)$ is a function on Σ_X solving the equation*

$$v(z) = I - \frac{1}{2\pi i} \int_{\Sigma_X} \frac{v(u)j_X^0(u)}{z_- - u} du \quad \text{for } z \in \Sigma_X \quad (11.8)$$

where z_- means the value of the integral on the minus side of Σ_X . Then

$$\mathbf{X}_n(z) = I - \frac{1}{2\pi i} \int_{\Sigma_X} \frac{v(u)j_X^0(u)}{z - u} du \quad \text{for } z \in \mathbb{C} \setminus \Sigma_X \quad (11.9)$$

solves the RHP for $\mathbf{X}_n$.

The proof of this lemma is immediate from the jump property of the Cauchy transform. By assumption

$$\mathbf{X}_{n-}(z) = v(z) \quad (11.10)$$

and the additive jump of the Cauchy transform gives

$$\mathbf{X}_{n+}(z) - \mathbf{X}_{n-}(z) = v(z)j_X^0(z) = \mathbf{X}_{n-}(z)j_X^0(z), \quad (11.11)$$

thus $\mathbf{X}_{n+}(z) = \mathbf{X}_{n-}(z)j_X(z)$. Asymptotics at infinity are given by (11.9).

The solution to equation (11.8) is given by a series of perturbation theory. Namely, the solution is

$$v(z) = I + \sum_{k=1}^{\infty} v_k(z) \quad (11.12)$$

where

$$v_k(z) = -\frac{1}{2\pi i} \int_{\Sigma_X} \frac{v_{k-1}(u)j_X^0(u)}{z - u} du, \quad v_0(z) = I. \quad (11.13)$$

This function clearly solves (11.8) provided the series converges, which it does, for sufficiently large n . Indeed, by (11.5),

$$|v_k(z)| \leq \left(\frac{C}{n}\right)^k \frac{1}{1 + |z|} \quad \text{for some constant } C > 0 \quad (11.14)$$

thus the series (11.12) is dominated by a convergent geometric series and thus converges absolutely. This in turn gives

$$\mathbf{X}_n(z) = I + \sum_{k=1}^{\infty} \mathbf{X}_{n,k}(z) \quad (11.15)$$

where

$$\mathbf{X}_{n,k}(z) = -\frac{1}{2\pi i} \int_{\Sigma_X} \frac{v_{k-1}(u) j_X^0(u)}{z-u} du. \quad (11.16)$$

We will need to compute

$$\mathbf{X}_{n,1}(z) = -\frac{1}{2\pi i} \int_{\Sigma_X} \frac{j_X^0(u)}{z-u} du. \quad (11.17)$$

12. EVALUATION OF $\mathbf{X}_1$

We are interested in the matrix $\mathbf{X}_1$, which gives the $\frac{1}{z}$ -term of $\mathbf{X}_n(z)$ at infinity, see (11.4). By (11.9),

$$\mathbf{X}_1 = -\frac{1}{2\pi i} \int_{\Sigma_X} v(u) j_X^0(u) du, \quad (12.1)$$

hence by (11.12), (11.14),

$$\mathbf{X}_1 = -\frac{1}{2\pi i} \int_{\Sigma_X} j_X^0(u) du + O(n^{-2}). \quad (12.2)$$

We would like to evaluate the integral,

$$-\frac{1}{2\pi i} \int_{\Sigma_X} j_X^0(u) du \quad (12.3)$$

with an error of the order of n^{-2} . By (11.7), it is enough to evaluate this integral over the circles $\partial D(\alpha, \varepsilon)$, $\partial D(\alpha', \varepsilon)$, $\partial D(\beta', \varepsilon)$, and $\partial D(\beta, \varepsilon)$. As we will see in the next section, the matrix-valued function $j_X^0(z)$ is analytic in the punctured disks, hence

$$\mathbf{X}_1 = -\left(\operatorname{Res}_{z=\alpha} + \operatorname{Res}_{z=\alpha'} + \operatorname{Res}_{z=\beta'} + \operatorname{Res}_{z=\beta} \right) j_X^0(z) + O(n^{-2}). \quad (12.4)$$

We will be especially interested in evaluation of the [12] element of the matrix $\mathbf{X}_1$, and we will prove the following asymptotic formula. Introduce the numbers,

$$\begin{aligned}\eta_\alpha &= \left[5 \frac{\vartheta_4''(n\omega + \frac{\omega}{2})}{\vartheta_4(n\omega + \frac{\omega}{2})} - 5 \frac{\vartheta_3''(\frac{\omega}{2})}{\vartheta_3(\frac{\omega}{2})} + 7 \left(\frac{\vartheta_4'(n\omega + \frac{\omega}{2})}{\vartheta_4(n\omega + \frac{\omega}{2})} \right)^2 + 17 \left(\frac{\vartheta_3'(\frac{\omega}{2})}{\vartheta_3(\frac{\omega}{2})} \right)^2 - 24 \frac{\vartheta_4'(n\omega + \frac{\omega}{2})\vartheta_3'(\frac{\omega}{2})}{\vartheta_4(n\omega + \frac{\omega}{2})\vartheta_3(\frac{\omega}{2})} \right], \\ \eta_{\alpha'} &= - \left[5 \frac{\vartheta_1''(n\omega + \frac{\omega}{2})}{\vartheta_1(n\omega + \frac{\omega}{2})} - 5 \frac{\vartheta_2''(\frac{\omega}{2})}{\vartheta_2(\frac{\omega}{2})} + 7 \left(\frac{\vartheta_1'(n\omega + \frac{\omega}{2})}{\vartheta_1(n\omega + \frac{\omega}{2})} \right)^2 + 17 \left(\frac{\vartheta_2'(\frac{\omega}{2})}{\vartheta_2(\frac{\omega}{2})} \right)^2 - 24 \frac{\vartheta_1'(n\omega + \frac{\omega}{2})\vartheta_2'(\frac{\omega}{2})}{\vartheta_1(n\omega + \frac{\omega}{2})\vartheta_2(\frac{\omega}{2})} \right], \\ \eta_{\beta'} &= - \left[5 \frac{\vartheta_2''(n\omega + \frac{\omega}{2})}{\vartheta_2(n\omega + \frac{\omega}{2})} - 5 \frac{\vartheta_1''(\frac{\omega}{2})}{\vartheta_1(\frac{\omega}{2})} + 7 \left(\frac{\vartheta_2'(n\omega + \frac{\omega}{2})}{\vartheta_2(n\omega + \frac{\omega}{2})} \right)^2 + 17 \left(\frac{\vartheta_1'(\frac{\omega}{2})}{\vartheta_1(\frac{\omega}{2})} \right)^2 - 24 \frac{\vartheta_2'(n\omega + \frac{\omega}{2})\vartheta_1'(\frac{\omega}{2})}{\vartheta_2(n\omega + \frac{\omega}{2})\vartheta_1(\frac{\omega}{2})} \right], \\ \eta_\beta &= \left[5 \frac{\vartheta_3''(n\omega + \frac{\omega}{2})}{\vartheta_3(n\omega + \frac{\omega}{2})} - 5 \frac{\vartheta_4''(\frac{\omega}{2})}{\vartheta_4(\frac{\omega}{2})} + 7 \left(\frac{\vartheta_3'(n\omega + \frac{\omega}{2})}{\vartheta_3(n\omega + \frac{\omega}{2})} \right)^2 + 17 \left(\frac{\vartheta_4'(\frac{\omega}{2})}{\vartheta_4(\frac{\omega}{2})} \right)^2 - 24 \frac{\vartheta_3'(n\omega + \frac{\omega}{2})\vartheta_4'(\frac{\omega}{2})}{\vartheta_3(n\omega + \frac{\omega}{2})\vartheta_4(\frac{\omega}{2})} \right],\end{aligned}\tag{12.5}$$

and

$$\begin{aligned}C_\alpha &= \frac{7}{2}(\beta' - \alpha) + \frac{3}{2}(\beta - \alpha) + \frac{3}{2}(\alpha' - \alpha) - \frac{(\alpha' - \alpha)(\beta - \alpha)}{(\beta' - \alpha)}, \\ C_{\alpha'} &= -\frac{7}{2}(\beta - \alpha') - \frac{3}{2}(\beta' - \alpha') + \frac{3}{2}(\alpha' - \alpha) - \frac{(\alpha' - \alpha)(\beta' - \alpha')}{(\beta - \alpha')}, \\ C_{\beta'} &= -\frac{7}{2}(\beta' - \alpha) - \frac{3}{2}(\beta' - \alpha') + \frac{3}{2}(\beta - \beta') - \frac{(\beta - \beta')(\beta' - \alpha')}{(\beta' - \alpha)}, \\ C_\beta &= \frac{7}{2}(\beta - \alpha') + \frac{3}{2}(\beta - \alpha) + \frac{3}{2}(\beta - \beta') - \frac{(\beta - \beta')(\beta - \alpha)}{(\beta - \alpha')}.\end{aligned}\tag{12.6}$$

Introduce also the numbers,

$$\begin{aligned}\Xi_\alpha &= \frac{\vartheta_3^2(0)\vartheta_4^2(n\omega + \frac{\omega}{2})}{\vartheta_3^2(\frac{\omega}{2})\vartheta_4^2(n\omega)}, & \Xi_{\alpha'} &= \frac{\vartheta_3^2(0)\vartheta_1^2(n\omega + \frac{\omega}{2})}{\vartheta_2^2(\frac{\omega}{2})\vartheta_4^2(n\omega)}, \\ \Xi_{\beta'} &= \frac{\vartheta_3^2(0)\vartheta_2^2(n\omega + \frac{\omega}{2})}{\vartheta_1^2(\frac{\omega}{2})\vartheta_4^2(n\omega)}, & \Xi_\beta &= \frac{\vartheta_3^2(0)\vartheta_3^2(n\omega + \frac{\omega}{2})}{\vartheta_4^2(\frac{\omega}{2})\vartheta_4^2(n\omega)},\end{aligned}\tag{12.7}$$

and

$$\begin{aligned}\xi_\alpha &= \frac{\vartheta_3'(\frac{\omega}{2})}{\vartheta_3(\frac{\omega}{2})} - \frac{\vartheta_4'(n\omega + \frac{\omega}{2})}{\vartheta_4(n\omega + \frac{\omega}{2})}, & \xi_{\alpha'} &= - \left(\frac{\vartheta_2'(\frac{\omega}{2})}{\vartheta_2(\frac{\omega}{2})} - \frac{\vartheta_1'(n\omega + \frac{\omega}{2})}{\vartheta_1(n\omega + \frac{\omega}{2})} \right), \\ \xi_{\beta'} &= \frac{\vartheta_1'(\frac{\omega}{2})}{\vartheta_1(\frac{\omega}{2})} - \frac{\vartheta_2'(n\omega + \frac{\omega}{2})}{\vartheta_2(n\omega + \frac{\omega}{2})}, & \xi_\beta &= - \left(\frac{\vartheta_4'(\frac{\omega}{2})}{\vartheta_4(\frac{\omega}{2})} - \frac{\vartheta_3'(n\omega + \frac{\omega}{2})}{\vartheta_3(n\omega + \frac{\omega}{2})} \right).\end{aligned}\tag{12.8}$$

Lemma 12.1. *As $n \rightarrow \infty$,*

$$[\mathbf{X}_1]_{12} = \frac{1}{n} (X_\alpha + X_{\alpha'} + X_{\beta'} + X_\beta) + O(n^{-2}),\tag{12.9}$$

where

$$\begin{aligned}
X_\alpha &= \frac{i\Xi_\alpha}{96} \left(C_\alpha + 12\pi\xi_\alpha + \frac{\pi^2\eta_\alpha}{2(\beta' - \alpha)} \right), \\
X_{\alpha'} &= \frac{i\Xi'_{\alpha'}}{96} \left(C_{\alpha'} + 12\pi\xi_{\alpha'} + \frac{\pi^2\eta_{\alpha'}}{2(\beta - \alpha')} \right), \\
X_{\beta'} &= \frac{i\Xi_{\beta'}}{96} \left(C_{\beta'} + 12\pi\xi_{\beta'} + \frac{\pi^2\eta_{\beta'}}{2(\beta' - \alpha)} \right), \\
X_\beta &= \frac{i\Xi_\beta}{96} \left(C_\beta + 12\pi\xi_\beta + \frac{\pi^2\eta_\beta}{2(\beta - \alpha')} \right).
\end{aligned} \tag{12.10}$$

Proof of this lemma is given in the next section.

13. PROOF OF LEMMA 12.1

On the circles, $\partial D(\alpha, \varepsilon)$ and $\partial D(\beta, \varepsilon)$, we have

$$j_X(z) = \begin{cases} = I - \frac{\psi_\alpha(z)^{-3/2}}{48n} \mathbf{M}(z) \begin{pmatrix} 1 & -6i \\ -6i & -1 \end{pmatrix} \mathbf{M}^{-1}(z) + O(n^{-2}) & \text{for } z \in \partial D(\alpha, \varepsilon), \\ = I - \frac{\psi_\beta(z)^{-3/2}}{48n} \mathbf{M}(z) \begin{pmatrix} 1 & 6i \\ 6i & -1 \end{pmatrix} \mathbf{M}^{-1}(z) + O(n^{-2}) & \text{for } z \in \partial D(\beta, \varepsilon), \end{cases} \tag{13.1}$$

and on the circles, $\partial D(\alpha', \varepsilon)$ and $\partial D(\beta', \varepsilon)$, we have

$$j_X(z) = \begin{cases} I - \frac{\psi_{\alpha'}(z)^{-3/2}}{48n} \mathbf{M}(z) \begin{pmatrix} -1 & -6ie^{\pm i\Omega_n} \\ -6ie^{\mp i\Omega_n} & 1 \end{pmatrix} \mathbf{M}^{-1}(z) + O(n^{-2}) \\ \quad \text{for } z \in \partial D(\alpha', \varepsilon), \pm \text{Im } z > 0, \\ I - \frac{\psi_{\beta'}(z)^{-3/2}}{48n} \mathbf{M}(z) \begin{pmatrix} -1 & 6ie^{\pm i\Omega_n} \\ 6ie^{\mp i\Omega_n} & 1 \end{pmatrix} \mathbf{M}^{-1}(z) + O(n^{-2}) \\ \quad \text{for } z \in \partial D(\beta', \varepsilon), \pm \text{Im } z > 0. \end{cases} \tag{13.2}$$

Thus

$$j_X^0(z) = \begin{cases} -\frac{\psi_\alpha(z)^{-3/2}}{48n} \mathbf{M}(z) \begin{pmatrix} 1 & -6i \\ -6i & -1 \end{pmatrix} \mathbf{M}^{-1}(z) + O(n^{-2}) & \text{for } z \in \partial D(\alpha, \varepsilon), \\ -\frac{\psi_\beta(z)^{-3/2}}{48n} \mathbf{M}(z) \begin{pmatrix} 1 & 6i \\ 6i & -1 \end{pmatrix} \mathbf{M}^{-1}(z) + O(n^{-2}) & \text{for } z \in \partial D(\beta, \varepsilon), \\ -\frac{\psi_{\alpha'}(z)^{-3/2}}{48n} \mathbf{M}(z) \begin{pmatrix} -1 & -6ie^{\pm i\Omega_n} \\ -6ie^{\mp i\Omega_n} & 1 \end{pmatrix} \mathbf{M}^{-1}(z) + O(n^{-2}) & \text{for } z \in \partial D(\alpha', \varepsilon), \\ -\frac{\psi_{\beta'}(z)^{-3/2}}{48n} \mathbf{M}(z) \begin{pmatrix} -1 & 6ie^{\pm i\Omega_n} \\ 6ie^{\mp i\Omega_n} & 1 \end{pmatrix} \mathbf{M}^{-1}(z) + O(n^{-2}) & \text{for } z \in \partial D(\beta', \varepsilon), \end{cases} \tag{13.3}$$

for $\pm \text{Im } z > 0$. To simplify notation, we will write the model solution given in (8.29) and (8.30) as

$$\mathbf{M}(z) = \frac{1}{2} \begin{pmatrix} (\gamma(z) + \gamma^{-1}(z))\vartheta_{11}(z) & i(\gamma(z) - \gamma^{-1}(z))\vartheta_{12}(z) \\ -i(\gamma(z) - \gamma^{-1}(z))\vartheta_{21}(z) & (\gamma(z) + \gamma^{-1}(z))\vartheta_{22}(z) \end{pmatrix} \quad (13.4)$$

where

$$\begin{aligned} \vartheta_{11}(z) &= \frac{\vartheta_3(\tilde{u}(z) - \tilde{u}_\infty + \frac{\Omega_n}{2})\vartheta_3(0)}{\vartheta_3(\tilde{u}(z) - \tilde{u}_\infty)\vartheta_3(\frac{\Omega_n}{2})} & \vartheta_{12}(z) &= \frac{\vartheta_3(\tilde{u}(z) + \tilde{u}_\infty - \frac{\Omega_n}{2})\vartheta_3(0)}{\vartheta_3(\tilde{u}(z) + \tilde{u}_\infty)\vartheta_3(\frac{\Omega_n}{2})} \\ \vartheta_{21}(z) &= \frac{\vartheta_3(\tilde{u}(z) + \tilde{u}_\infty + \frac{\Omega_n}{2})\vartheta_3(0)}{\vartheta_3(\tilde{u}(z) + \tilde{u}_\infty)\vartheta_3(\frac{\Omega_n}{2})} & \vartheta_{22}(z) &= \frac{\vartheta_3(\tilde{u}(z) - \tilde{u}_\infty - \frac{\Omega_n}{2})\vartheta_3(0)}{\vartheta_3(\tilde{u}(z) - \tilde{u}_\infty)\vartheta_3(\frac{\Omega_n}{2})}. \end{aligned} \quad (13.5)$$

Notice that each of the functions ϑ_{ij} is analytic throughout the complex plane, except on the intervals (α, α') and (β', β) , where they satisfy the relations

$$[\vartheta_{11}]_\pm = [\vartheta_{12}]_\mp, \quad [\vartheta_{21}]_\pm = [\vartheta_{22}]_\mp, \quad (13.6)$$

and on the interval (α', β') , where they satisfy

$$[\vartheta_{11}]_+ = e^{-i\Omega_n}[\vartheta_{11}]_-, \quad [\vartheta_{12}]_+ = e^{i\Omega_n}[\vartheta_{12}]_-, \quad [\vartheta_{21}]_+ = e^{-i\Omega_n}[\vartheta_{21}]_-, \quad [\vartheta_{22}]_+ = e^{i\Omega_n}[\vartheta_{22}]_-. \quad (13.7)$$

Multiplying out equations (13.3) gives

$$j_X^0(z) = \frac{\psi_\xi(z)^{-3/2}}{48n} \begin{pmatrix} j_{11}^\xi & j_{12}^\xi \\ j_{21}^\xi & j_{22}^\xi \end{pmatrix} + O(n^{-2}) \quad \text{for } z \in \partial D(\xi, \varepsilon), \quad \xi = \alpha, \alpha', \beta', \beta, \quad (13.8)$$

where

$$\begin{aligned} j_{12}^\alpha &= \frac{i}{2} [3((\gamma^2(z) + \gamma^{-2}(z))(\vartheta_{11}^2 - \vartheta_{12}^2) + (\gamma^2(z) - \gamma^{-2}(z))\vartheta_{11}\vartheta_{12} + 6(\vartheta_{11}^2 - \vartheta_{12}^2))], \\ j_{12}^{\alpha'} &= \frac{i}{2} [3(\gamma^2(z) + \gamma^{-2}(z))J(z) - (\gamma^2(z) - \gamma^{-2}(z))\vartheta_{11}\vartheta_{12} + 6K(z)], \\ j_{12}^{\beta'} &= -\frac{i}{2} [3(\gamma^2(z) + \gamma^{-2}(z))J(z) + (\gamma^2(z) - \gamma^{-2}(z))\vartheta_{11}\vartheta_{12} + 6K(z)], \\ j_{12}^\beta &= -\frac{i}{2} [3((\gamma^2(z) + \gamma^{-2}(z))(\vartheta_{11}^2 - \vartheta_{12}^2) - (\gamma^2(z) - \gamma^{-2}(z))\vartheta_{11}\vartheta_{12} + 6(\vartheta_{11}^2 - \vartheta_{12}^2))], \end{aligned} \quad (13.9)$$

and

$$J(z) = \begin{cases} \vartheta_{11}^2 e^{i\Omega_n} + \vartheta_{12}^2 e^{-i\Omega_n} & \text{for } \text{Im } z > 0, \\ \vartheta_{11}^2 e^{-i\Omega_n} + \vartheta_{12}^2 e^{i\Omega_n} & \text{for } \text{Im } z < 0, \end{cases} \quad (13.10)$$

$$K(z) = \begin{cases} \vartheta_{11}^2 e^{i\Omega_n} - \vartheta_{12}^2 e^{-i\Omega_n} & \text{for } \text{Im } z > 0, \\ \vartheta_{11}^2 e^{-i\Omega_n} - \vartheta_{12}^2 e^{i\Omega_n} & \text{for } \text{Im } z < 0. \end{cases} \quad (13.11)$$

In order to integrate $j_X^0(z)$, let us examine the behavior of the various functions described above near each of the turning points. Introduce the numbers

$$\begin{aligned} A_\alpha &= \sqrt{(\alpha' - \alpha)(\beta' - \alpha)(\beta - \alpha)}, & A_{\alpha'} &= \sqrt{(\alpha' - \alpha)(\beta' - \alpha')(\beta - \alpha')}, \\ A_{\beta'} &= \sqrt{(\beta' - \alpha)(\beta' - \alpha')(\beta - \beta')}, & A_\beta &= \sqrt{(\beta - \alpha)(\beta - \alpha')(\beta - \beta')}, \end{aligned} \quad (13.12)$$

and

$$\begin{aligned} B_\alpha &= \frac{1}{\alpha' - \alpha} + \frac{1}{\beta' - \alpha} + \frac{1}{\beta - \alpha}, & B_{\alpha'} &= -\frac{1}{\alpha' - \alpha} + \frac{1}{\beta' - \alpha'} + \frac{1}{\beta - \alpha'}, \\ B_{\beta'} &= \frac{1}{\beta' - \alpha} + \frac{1}{\beta' - \alpha'} - \frac{1}{\beta - \beta'}, & B_\beta &= \frac{1}{\beta - \alpha} + \frac{1}{\beta - \alpha'} + \frac{1}{\beta - \beta'}. \end{aligned} \quad (13.13)$$

For $x \in (\beta, \beta + \varepsilon)$, we have

$$\tilde{u}(x) = \frac{\pi}{2K}u(x) = \frac{\pi}{A_\beta}\sqrt{x - \beta} + O((x - \beta)^{3/2}); \quad (13.14)$$

for $x \in (\alpha - \varepsilon, \alpha)$,

$$\tilde{u}(x) = \frac{\pi}{2K}u(x) = \frac{\pi}{2} - \frac{\pi}{A_\alpha}\sqrt{\alpha - x} + O((\alpha - x)^{3/2}); \quad (13.15)$$

for $x \in (\alpha', \alpha' + \varepsilon)$,

$$\tilde{u}_\pm(x) = \frac{\pi}{2K}u_\pm(x) = \frac{\pi}{2} \pm \frac{\tau\pi}{2} - \frac{\pi}{A_{\alpha'}}\sqrt{x - \alpha'} + O((x - \alpha')^{3/2}); \quad (13.16)$$

and for $x \in (\beta' - \varepsilon, \beta')$,

$$\tilde{u}_\pm(x) = \frac{\pi}{2K}u_\pm(x) = \pm \frac{\tau\pi}{2} + \frac{\pi}{A_{\beta'}}\sqrt{\beta' - x} + O((\beta' - x)^{3/2}). \quad (13.17)$$

Also, from (8.4) we have that

$$\gamma^2(z) \pm \gamma^{-2}(z) = \sqrt{\frac{(z - \alpha)(z - \beta')}{(z - \alpha')(z - \beta)}} \pm \sqrt{\frac{(z - \alpha')(z - \beta)}{(z - \alpha)(z - \beta')}}, \quad (13.18)$$

and from (9.21), (9.28), (10.9), (10.27), and (2.19) that

$$\begin{aligned} \psi_\alpha^{-3/2}(z) &= (\alpha - z)^{-3/2} \left[\frac{A_\alpha}{2} + \frac{1}{20}A_\alpha B_\alpha(\alpha - z) + O((\alpha - z)^2) \right], \\ \psi_{\alpha'}^{-3/2}(z) &= (z - \alpha')^{-3/2} \left[\frac{A_{\alpha'}}{2} - \frac{1}{20}A_{\alpha'} B_{\alpha'}(z - \alpha') + O((z - \alpha')^2) \right], \\ \psi_{\beta'}^{-3/2}(z) &= (\beta' - z)^{-3/2} \left[\frac{A_{\beta'}}{2} - \frac{1}{20}A_{\beta'} B_{\beta'}(\beta' - z) + O((\beta' - z)^2) \right], \\ \psi_\beta^{-3/2}(z) &= (z - \beta)^{-3/2} \left[\frac{A_\beta}{2} + \frac{1}{20}A_\beta B_\beta(z - \beta) + O((z - \beta)^2) \right]. \end{aligned} \quad (13.19)$$

It follows that the functions $(\gamma^2 \pm \gamma^{-2})\psi^{-3/2}(z)$ are meromorphic in a neighborhood of each of the turning points. In particular, at $z = \alpha$, we have

$$\begin{aligned} (\gamma^2 \pm \gamma^{-2})\psi_\alpha^{-3/2}(z) &= \pm \frac{(\alpha' - \alpha)(\beta - \alpha)}{2(\alpha - z)^2} + \frac{1}{(\alpha - z)} \left[\frac{(\beta' - \alpha)}{2} \pm \frac{3}{10}(\beta - \alpha) \right. \\ &\quad \left. \pm \frac{3}{10}(\alpha' - \alpha) \mp \frac{1}{5} \frac{(\alpha' - \alpha)(\beta - \alpha)}{(\beta' - \alpha)} \right] + O(1); \end{aligned} \quad (13.20)$$

at $z = \beta$, we have

$$\begin{aligned} (\gamma^2 \pm \gamma^{-2})\psi_\beta^{-3/2}(z) &= \frac{(\beta - \alpha)(\beta - \beta')}{2(z - \beta)^2} + \frac{1}{(z - \beta)} \left[\pm \frac{1}{2}(\beta - \alpha') + \frac{3}{10}(\beta - \alpha) \right. \\ &\quad \left. + \frac{3}{10}(\beta - \beta') - \frac{1}{5} \frac{(\beta - \alpha)(\beta - \beta')}{(\beta - \alpha')} \right] + O(1); \end{aligned} \quad (13.21)$$

at $z = \alpha'$, we have

$$\begin{aligned} (\gamma^2 \pm \gamma^{-2})\psi_{\alpha'}^{-3/2}(z) &= \frac{(\alpha' - \alpha)(\beta' - \alpha')}{2(z - \alpha')^2} + \frac{1}{(z - \alpha')} \left[\pm \frac{(\beta - \alpha')}{2} + \frac{3}{10}(\beta' - \alpha') \right. \\ &\quad \left. - \frac{3}{10}(\alpha' - \alpha) + \frac{1}{5} \frac{(\alpha' - \alpha)(\beta' - \alpha')}{(\beta - \alpha')} \right] + O(1); \end{aligned} \quad (13.22)$$

and at $z = \beta'$, we have

$$\begin{aligned} (\gamma^2 \pm \gamma^{-2})\psi_{\beta'}^{-3/2}(z) &= \pm \frac{(\beta' - \alpha')(\beta - \beta')}{2(\beta' - z)^2} + \frac{1}{(\beta' - z)} \left[\frac{(\beta' - \alpha)}{2} \pm \frac{3}{10}(\beta' - \alpha') \right. \\ &\quad \left. \mp \frac{3}{10}(\beta - \beta') \pm \frac{1}{5} \frac{(\beta' - \alpha')(\beta - \beta')}{\beta' - \alpha} \right] + O(1). \end{aligned} \quad (13.23)$$

Notice also, from the relations (13.5), that the functions $\vartheta_{11}^2 + \vartheta_{12}^2$ and $\vartheta_{11}\vartheta_{12}$ have no jumps in neighborhoods of α or β , and take finite values at $z = \alpha$ and $z = \beta$, thus are analytic in neighborhoods of α and β . Using (13.14) and (13.15), we see that these functions have Taylor expansions about $z = \beta$,

$$\begin{aligned} \vartheta_{11}^2(z) + \vartheta_{12}^2(z) &= 2 \frac{\vartheta_3^2(0)}{\vartheta_3^2(\frac{\Omega_n}{2})} \frac{\vartheta_3^2(\tilde{u}_\infty - \frac{\Omega_n}{2})}{\vartheta_3^2(\tilde{u}_\infty)} + \frac{\pi^2}{2A_\beta^2} \frac{d^2}{d\tilde{u}^2} \left(\vartheta_{11}^2 + \vartheta_{12}^2 \right) \Big|_{z=\beta} (z - \beta) + \dots, \\ \vartheta_{11}(z)\vartheta_{12}(z) &= \frac{\vartheta_3^2(0)}{\vartheta_3^2(\frac{\Omega_n}{2})} \frac{\vartheta_3^2(\tilde{u}_\infty - \frac{\Omega_n}{2})}{\vartheta_3^2(\tilde{u}_\infty)} + \frac{\pi^2}{2A_\beta^2} \frac{d^2}{d\tilde{u}^2} \left(\vartheta_{11}\vartheta_{12} \right) \Big|_{z=\beta} (z - \beta) + \dots, \end{aligned} \quad (13.24)$$

and about $z = \alpha$,

$$\begin{aligned} \vartheta_{11}^2(z) + \vartheta_{12}^2(z) &= 2 \frac{\vartheta_3^2(0)}{\vartheta_3^2(\frac{\Omega_n}{2})} \frac{\vartheta_4^2(\tilde{u}_\infty - \frac{\Omega_n}{2})}{\vartheta_4^2(\tilde{u}_\infty)} + \frac{\pi^2}{2A_\alpha^2} \frac{d^2}{d\tilde{u}^2} \left(\vartheta_{11}^2 + \vartheta_{12}^2 \right) \Big|_{z=\alpha} (\alpha - z) + \dots, \\ \vartheta_{11}(z)\vartheta_{12}(z) &= \frac{\vartheta_3^2(0)}{\vartheta_3^2(\frac{\Omega_n}{2})} \frac{\vartheta_4^2(\tilde{u}_\infty - \frac{\Omega_n}{2})}{\vartheta_4^2(\tilde{u}_\infty)} + \frac{\pi^2}{2A_\alpha^2} \frac{d^2}{d\tilde{u}^2} \left(\vartheta_{11}\vartheta_{12} \right) \Big|_{z=\alpha} (\alpha - z) + \dots. \end{aligned} \quad (13.25)$$

By a similar argument, $J(z)$ and $\vartheta_{11}\vartheta_{12}$ are also analytic in neighborhoods of α' and β' and using (13.16) and (13.17) we can write their Taylor expansions about $z = \alpha'$,

$$\begin{aligned} J(z) &= 2 \frac{\vartheta_3^2(0)}{\vartheta_3^2(\frac{\Omega_n}{2})} \frac{\vartheta_1^2(\tilde{u}_\infty - \frac{\Omega_n}{2})}{\vartheta_1^2(\tilde{u}_\infty)} + \frac{\pi^2}{2A_{\alpha'}^2} \frac{d^2}{d\tilde{u}^2} J(z) \Big|_{z=\alpha'} (z - \alpha') + \dots, \\ \vartheta_{11}(z)\vartheta_{12}(z) &= \frac{\vartheta_3^2(0)}{\vartheta_3^2(\frac{\Omega_n}{2})} \frac{\vartheta_1^2(\tilde{u}_\infty - \frac{\Omega_n}{2})}{\vartheta_1^2(\tilde{u}_\infty)} + \frac{\pi^2}{2A_{\alpha'}^2} \frac{d^2}{d\tilde{u}^2} \left(\vartheta_{11}\vartheta_{12} \right) \Big|_{z=\alpha'} (z - \alpha') + \dots, \end{aligned} \quad (13.26)$$

and about $z = \beta'$,

$$\begin{aligned} J(z) &= 2 \frac{\vartheta_3^2(0)}{\vartheta_3^2(\frac{\Omega_n}{2})} \frac{\vartheta_2^2(\tilde{u}_\infty - \frac{\Omega_n}{2})}{\vartheta_2^2(\tilde{u}_\infty)} + \frac{\pi^2}{2A_{\beta'}^2} \frac{d^2}{d\tilde{u}^2} J(z) \Big|_{z=\beta'} (\beta' - z) + \dots, \\ \vartheta_{11}(z)\vartheta_{12}(z) &= \frac{\vartheta_3^2(0)}{\vartheta_3^2(\frac{\Omega_n}{2})} \frac{\vartheta_2^2(\tilde{u}_\infty - \frac{\Omega_n}{2})}{\vartheta_2^2(\tilde{u}_\infty)} + \frac{\pi^2}{2A_{\beta'}^2} \frac{d^2}{d\tilde{u}^2} \left(\vartheta_{11}\vartheta_{12} \right) \Big|_{z=\beta'} (\beta' - z) + \dots. \end{aligned} \quad (13.27)$$

Finally, notice that the function $\vartheta_{11}^2 - \vartheta_{12}^2$ is an odd function of $\tilde{u}$, and using (13.14) and (13.15), we can write, for $x \in (\beta, \beta + \varepsilon)$,

$$\begin{aligned} \vartheta_{11}^2(x) - \vartheta_{12}^2(x) &= \sqrt{x - \beta} \\ &\times \left[\frac{4\pi}{A_\beta} \frac{\vartheta_3^2(0)}{\vartheta_3^2(\frac{\Omega_n}{2})} \frac{\vartheta_3^2(\tilde{u}_\infty - \frac{\Omega_n}{2})}{\vartheta_3^2(\tilde{u}_\infty)} \left(\frac{\vartheta_3'(\tilde{u}_\infty)}{\vartheta_3(\tilde{u}_\infty)} - \frac{\vartheta_3'(\tilde{u}_\infty - \frac{\Omega_n}{2})}{\vartheta_3(\tilde{u}_\infty - \frac{\Omega_n}{2})} \right) + O(x - \beta) \right], \end{aligned} \quad (13.28)$$

and for $x \in (\alpha - \varepsilon, \alpha)$,

$$\begin{aligned} \vartheta_{11}^2(x) - \vartheta_{12}^2(x) &= -\sqrt{\alpha - x} \\ &\times \left[\frac{4\pi}{A_\alpha} \frac{\vartheta_3^2(0)}{\vartheta_3^2(\frac{\Omega_n}{2})} \frac{\vartheta_4^2(\tilde{u}_\infty - \frac{\Omega_n}{2})}{\vartheta_4^2(\tilde{u}_\infty)} \left(\frac{\vartheta_4'(\tilde{u}_\infty)}{\vartheta_4(\tilde{u}_\infty)} - \frac{\vartheta_4'(\tilde{u}_\infty - \frac{\Omega_n}{2})}{\vartheta_4(\tilde{u}_\infty - \frac{\Omega_n}{2})} \right) + O(\alpha - x) \right]. \end{aligned} \quad (13.29)$$

Similarly, using (13.16) and (13.17), we can write, for $x \in (\alpha', \alpha' + \varepsilon)$,

$$\begin{aligned} K(x) &= -\sqrt{x - \alpha'} \\ &\times \left[\frac{4\pi}{A_{\alpha'}} \frac{\vartheta_3^2(0)}{\vartheta_3^2(\frac{\Omega_n}{2})} \frac{\vartheta_1^2(\tilde{u}_\infty - \frac{\Omega_n}{2})}{\vartheta_1^2(\tilde{u}_\infty)} \left(\frac{\vartheta_1'(\tilde{u}_\infty)}{\vartheta_1(\tilde{u}_\infty)} - \frac{\vartheta_1'(\tilde{u}_\infty - \frac{\Omega_n}{2})}{\vartheta_1(\tilde{u}_\infty - \frac{\Omega_n}{2})} \right) + O(x - \alpha') \right], \end{aligned} \quad (13.30)$$

and for $x \in (\beta' - \varepsilon, \beta')$,

$$\begin{aligned} K(x) &= \sqrt{\beta' - x} \\ &\times \left[\frac{4\pi}{A_{\beta'}} \frac{\vartheta_3^2(0)}{\vartheta_3^2(\frac{\Omega_n}{2})} \frac{\vartheta_2^2(\tilde{u}_\infty - \frac{\Omega_n}{2})}{\vartheta_2^2(\tilde{u}_\infty)} \left(\frac{\vartheta_2'(\tilde{u}_\infty)}{\vartheta_2(\tilde{u}_\infty)} - \frac{\vartheta_2'(\tilde{u}_\infty - \frac{\Omega_n}{2})}{\vartheta_2(\tilde{u}_\infty - \frac{\Omega_n}{2})} \right) + O(\beta' - x) \right] \end{aligned} \quad (13.31)$$

From equations (13.19), (13.28), (13.29), (13.30), and (13.31), it follows that the functions $(\vartheta_{11}^2(z) - \vartheta_{12}^2(z))\psi_\alpha^{-3/2}(z)$ and $(\vartheta_{11}^2(z) - \vartheta_{12}^2(z))\psi_\beta^{-3/2}(z)$ are meromorphic in neighborhoods of α and β , respectively, and have simple poles at $z = \alpha$ and $z = \beta$, respectively, and that the functions $K(z)\psi_{\alpha'}^{-3/2}(z)$ and $K(z)\psi_{\beta'}^{-3/2}(z)$ are meromorphic in neighborhoods of α' and β' , respectively, and have simple poles at $z = \alpha'$ and $z = \beta'$, respectively.

Let us compute the residues of functions that appear in (13.9). Observe that

$$\frac{\Omega_n}{2} = n\omega + \frac{\pi}{2}, \quad \tilde{u}_\infty = -\frac{\omega}{2} + \frac{\pi}{2}, \quad \frac{\Omega_n}{2} - \tilde{u}_\infty = n\omega + \frac{\omega}{2}. \quad (13.32)$$

From (13.20), (13.21), (13.24), and (13.25), we obtain that

$$\begin{aligned} & \text{Res}_{z=\alpha} 3(\vartheta_{11}^2(z) + \vartheta_{12}^2(z)) (\gamma^2(z) + \gamma^{-2}(z)) \psi_\alpha^{-3/2}(z) \\ &= \frac{\vartheta_3^2(0)\vartheta_4^2(n\omega + \frac{\omega}{2})}{\vartheta_3^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[-3(\beta' - \alpha) - \frac{9}{5}(\beta - \alpha) - \frac{9}{5}(\alpha' - \alpha) + \frac{6}{5} \frac{(\alpha' - \alpha)(\beta - \alpha)}{(\beta' - \alpha)} \right] \\ & \quad - \frac{3\pi^2}{4(\beta' - \alpha)} \frac{d^2}{d\tilde{u}^2} \left(\vartheta_{11}^2 + \vartheta_{12}^2 \right) \Big|_{z=\alpha} \end{aligned} \quad (13.33)$$

and

$$\begin{aligned} & \text{Res}_{z=\beta} 3(\vartheta_{11}^2(z) + \vartheta_{12}^2(z)) (\gamma^2(z) + \gamma^{-2}(z)) \psi_\beta^{-3/2}(z) \\ &= \frac{\vartheta_3^2(0)\vartheta_4^2(n\omega + \frac{\omega}{2})}{\vartheta_4^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[3(\beta - \alpha') + \frac{9}{5}(\beta - \beta') + \frac{9}{5}(\beta - \alpha) - \frac{6}{5} \frac{(\beta - \alpha)(\beta - \beta')}{(\beta - \alpha')} \right] \\ & \quad + \frac{3\pi^2}{4(\beta - \alpha')} \frac{d^2}{d\tilde{u}^2} \left(\vartheta_{11}^2 + \vartheta_{12}^2 \right) \Big|_{z=\beta}. \end{aligned} \quad (13.34)$$

Also,

$$\begin{aligned} & \text{Res}_{z=\alpha} (\vartheta_{11}(z)\vartheta_{12}(z)) (\gamma^2(z) - \gamma^{-2}(z)) \psi_\alpha^{-3/2}(z) \\ &= \frac{\vartheta_3^2(0)\vartheta_4^2(n\omega + \frac{\omega}{2})}{\vartheta_3^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[-\frac{(\beta' - \alpha)}{2} + \frac{3}{10}(\beta - \alpha) + \frac{3}{10}(\alpha' - \alpha) - \frac{1}{5} \frac{(\alpha' - \alpha)(\beta - \alpha)}{(\beta' - \alpha)} \right] \\ & \quad + \frac{\pi^2}{4(\beta' - \alpha)} \frac{d^2}{d\tilde{u}^2} \left(\vartheta_{11}\vartheta_{12} \right) \Big|_{z=\alpha}, \end{aligned} \quad (13.35)$$

and

$$\begin{aligned} & \text{Res}_{z=\beta} (\vartheta_{11}(z)\vartheta_{12}(z)) (\gamma^2(z) - \gamma^{-2}(z)) \psi_\beta^{-3/2}(z) \\ &= \frac{\vartheta_3^2(0)\vartheta_4^2(n\omega + \frac{\omega}{2})}{\vartheta_4^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[-\frac{(\beta - \alpha')}{2} + \frac{3}{10}(\beta - \beta') + \frac{3}{10}(\beta - \alpha) - \frac{1}{5} \frac{(\beta - \alpha)(\beta - \beta')}{(\beta - \alpha')} \right] \\ & \quad + \frac{\pi^2}{4(\beta - \alpha')} \frac{d^2}{d\tilde{u}^2} \left(\vartheta_{11}\vartheta_{12} \right) \Big|_{z=\beta}. \end{aligned} \quad (13.36)$$

From (13.19), (13.28), and (13.29), we obtain that

$$\text{Res}_{z=\alpha} 6(\vartheta_{11}^2(z) - \vartheta_{12}^2(z)) \psi_\alpha^{-3/2}(z) = -12\pi \frac{\vartheta_3^2(0)\vartheta_4^2(n\omega + \frac{\omega}{2})}{\vartheta_3^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[\frac{\vartheta_3'(\frac{\omega}{2})}{\vartheta_3(\frac{\omega}{2})} - \frac{\vartheta_4'(n\omega + \frac{\omega}{2})}{\vartheta_4(n\omega + \frac{\omega}{2})} \right] \quad (13.37)$$

and

$$\text{Res}_{z=\beta} 6(\vartheta_{11}^2(z) - \vartheta_{12}^2(z)) \psi_\beta^{-3/2}(z) = -12\pi \frac{\vartheta_3^2(0)\vartheta_4^2(n\omega + \frac{\omega}{2})}{\vartheta_4^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[\frac{\vartheta_4'(\frac{\omega}{2})}{\vartheta_4(\frac{\omega}{2})} - \frac{\vartheta_3'(n\omega + \frac{\omega}{2})}{\vartheta_3(n\omega + \frac{\omega}{2})} \right]. \quad (13.38)$$

We now turn our attention to the residues at the inner turning points. From (13.22), (13.23), (13.26), and (13.27), we have

$$\begin{aligned} & \text{Res}_{z=\alpha'} 3\psi_{\alpha'}^{-3/2}(z)J(z)(\gamma^2(z) + \gamma^{-2}(z)) \\ &= \frac{\vartheta_3^2(0)\vartheta_1^2(n\omega + \frac{\omega}{2})}{\vartheta_2^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[3(\beta - \alpha') + \frac{9}{5}(\beta' - \alpha') - \frac{9}{5}(\alpha' - \alpha) + \frac{6}{5} \frac{(\alpha' - \alpha)(\beta' - \alpha')}{\beta - \alpha'} \right] \\ & \quad + \frac{3\pi^2}{4(\beta - \alpha')} \frac{d^2}{d\tilde{u}^2} J(z) \Big|_{z=\alpha'} \end{aligned} \quad (13.39)$$

and

$$\begin{aligned} & \text{Res}_{z=\beta'} 3\psi_{\beta'}^{-3/2}(z)J(z)(\gamma^2(z) + \gamma^{-2}(z)) \\ &= \frac{\vartheta_3^2(0)\vartheta_2^2(n\omega + \frac{\omega}{2})}{\vartheta_1^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[-3(\beta' - \alpha) - \frac{9}{5}(\beta' - \alpha') + \frac{9}{5}(\beta - \beta') - \frac{6}{5} \frac{(\beta' - \alpha')(\beta - \beta')}{\beta' - \alpha} \right] \\ & \quad - \frac{3\pi^2}{4(\beta' - \alpha)} \frac{d^2}{d\tilde{u}^2} J(z) \Big|_{z=\beta'}. \end{aligned} \quad (13.40)$$

Also,

$$\begin{aligned} & \text{Res}_{z=\alpha'} \psi_{\alpha'}^{-3/2}(z)(\gamma^2(z) - \gamma^{-2}(z))\vartheta_{11}(z)\vartheta_{12}(z) \\ &= \frac{\vartheta_3^2(0)\vartheta_1^2(n\omega + \frac{\omega}{2})}{\vartheta_2^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[-\frac{(\beta - \alpha')}{2} + \frac{3}{10}(\beta' - \alpha') - \frac{3}{10}(\alpha' - \alpha) + \frac{1}{5} \frac{(\alpha' - \alpha)(\beta' - \alpha')}{\beta - \alpha'} \right] \\ & \quad + \frac{\pi^2}{4(\beta - \alpha')} \frac{d^2}{d\tilde{u}^2} \left(\vartheta_{11}\vartheta_{12} \right) \Big|_{z=\alpha'} \end{aligned} \quad (13.41)$$

and

$$\begin{aligned} & \text{Res}_{z=\beta'} \psi_{\beta'}^{-3/2}(z)(\gamma^2(z) - \gamma^{-2}(z))\vartheta_{11}(z)\vartheta_{12}(z) \\ &= \frac{\vartheta_3^2(0)\vartheta_2^2(n\omega + \frac{\omega}{2})}{\vartheta_1^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[-\frac{(\beta' - \alpha)}{2} + \frac{3}{10}(\beta' - \alpha') - \frac{3}{10}(\beta - \beta') + \frac{1}{5} \frac{(\beta' - \alpha')(\beta - \beta')}{\beta' - \alpha} \right] \\ & \quad + \frac{\pi^2}{4(\beta' - \alpha)} \frac{d^2}{d\tilde{u}^2} \left(\vartheta_{11}\vartheta_{12} \right) \Big|_{z=\beta'}. \end{aligned} \quad (13.42)$$

From (13.19), (13.30), (13.31), we have

$$\text{Res}_{z=\alpha'} 6\psi_{\alpha'}^{-3/2}(z)K(z) = 12\pi \frac{\vartheta_3^2(0)\vartheta_1^2(n\omega + \frac{\omega}{2})}{\vartheta_2^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[\frac{\vartheta_2'(\frac{\omega}{2})}{\vartheta_2(\frac{\omega}{2})} - \frac{\vartheta_1'(n\omega + \frac{\omega}{2})}{\vartheta_1(n\omega + \frac{\omega}{2})} \right], \quad (13.43)$$

and

$$\text{Res}_{z=\beta'} 6\psi_{\beta'}^{-3/2}(z)K(z) = 12\pi \frac{\vartheta_3^2(0)\vartheta_2^2(n\omega + \frac{\omega}{2})}{\vartheta_1^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[\frac{\vartheta_1'(\frac{\omega}{2})}{\vartheta_1(\frac{\omega}{2})} - \frac{\vartheta_2'(n\omega + \frac{\omega}{2})}{\vartheta_2(n\omega + \frac{\omega}{2})} \right]. \quad (13.44)$$

Combining (13.8),(13.9),(13.33),(13.35), and (13.37), we get that

$$\begin{aligned} \text{Res}_{z=\alpha} [j_X^0(z)] &= \frac{i}{96} \left[\frac{\vartheta_3^2(0)\vartheta_4^2(n\omega + \frac{\omega}{2})}{\vartheta_3^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left(-\frac{7}{2}(\beta' - \alpha) - \frac{3}{2}(\beta - \alpha) - \frac{3}{2}(\alpha' - \alpha) + \frac{(\alpha' - \alpha)(\beta - \alpha)}{\beta' - \alpha} \right) \right. \\ &\quad \left. - 12\pi \left(\frac{\vartheta'_3(\frac{\omega}{2})}{\vartheta_3(\frac{\omega}{2})} - \frac{\vartheta'_4(n\omega + \frac{\omega}{2})}{\vartheta_4(n\omega + \frac{\omega}{2})} \right) \right] - \frac{\pi^2}{4(\beta' - \alpha)} \frac{d^2}{d\tilde{u}^2} \left[3(\vartheta_{11}^2 + \vartheta_{12}^2) - \vartheta_{11}\vartheta_{12} \right]_{z=\alpha}. \end{aligned} \quad (13.45)$$

Similarly, we have

$$\begin{aligned} \text{Res}_{z=\alpha'} [j_X^0(z)] &= \frac{i}{96} \left[\frac{\vartheta_3^2(0)\vartheta_1^2(n\omega + \frac{\omega}{2})}{\vartheta_2^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left(\frac{7}{2}(\beta - \alpha') + \frac{3}{2}(\beta' - \alpha') - \frac{3}{2}(\alpha' - \alpha) + \frac{(\alpha' - \alpha)(\beta' - \alpha')}{\beta - \alpha'} \right) \right. \\ &\quad \left. + 12\pi \left(\frac{\vartheta'_2(\frac{\omega}{2})}{\vartheta_2(\frac{\omega}{2})} - \frac{\vartheta'_1(n\omega + \frac{\omega}{2})}{\vartheta_1(n\omega + \frac{\omega}{2})} \right) \right] + \frac{\pi^2}{4(\beta - \alpha')} \frac{d^2}{d\tilde{u}^2} \left[3J(z) - \vartheta_{11}\vartheta_{12} \right]_{z=\alpha'}, \end{aligned} \quad (13.46)$$

$$\begin{aligned} \text{Res}_{z=\beta'} [j_X^0(z)] &= \frac{i}{96} \left[\frac{\vartheta_3^2(0)\vartheta_2^2(n\omega + \frac{\omega}{2})}{\vartheta_1^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[\frac{7}{2}(\beta' - \alpha) + \frac{3}{2}(\beta' - \alpha') - \frac{3}{2}(\beta - \beta') + \frac{(\beta - \beta')(\beta' - \alpha')}{\beta' - \alpha} \right] \right. \\ &\quad \left. - 12\pi \left(\frac{\vartheta'_1(\frac{\omega}{2})}{\vartheta_1(\frac{\omega}{2})} - \frac{\vartheta'_2(n\omega + \frac{\omega}{2})}{\vartheta_2(n\omega + \frac{\omega}{2})} \right) \right] + \frac{\pi^2}{4(\beta' - \alpha)} \frac{d^2}{d\tilde{u}^2} \left[3J(z) - \vartheta_{11}\vartheta_{12} \right]_{z=\beta'}, \end{aligned} \quad (13.47)$$

and

$$\begin{aligned} \text{Res}_{z=\beta} [j_X^0(z)] &= \frac{i}{96} \left[\frac{\vartheta_3^2(0)\vartheta_3^2(n\omega + \frac{\omega}{2})}{\vartheta_4^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left(-\frac{7}{2}(\beta - \alpha') - \frac{3}{2}(\beta - \alpha) - \frac{3}{2}(\beta - \beta') + \frac{(\beta - \beta')(\beta - \alpha)}{\beta - \alpha'} \right) \right. \\ &\quad \left. + 12\pi \left(\frac{\vartheta'_4(\frac{\omega}{2})}{\vartheta_4(\frac{\omega}{2})} - \frac{\vartheta'_3(n\omega + \frac{\omega}{2})}{\vartheta_3(n\omega + \frac{\omega}{2})} \right) \right] - \frac{\pi^2}{4(\beta - \alpha')} \frac{d^2}{d\tilde{u}^2} \left[3(\vartheta_{11}^2 + \vartheta_{12}^2) - \vartheta_{11}\vartheta_{12} \right]_{z=\beta}. \end{aligned} \quad (13.48)$$

Using MAPLE for calculations, we get

$$\begin{aligned} \frac{d^2}{d\tilde{u}^2} [3(\vartheta_{11}^2 + \vartheta_{12}^2) - \vartheta_{11}\vartheta_{12}] \Big|_{z=\alpha} &= 2 \frac{\vartheta_3^2(0)\vartheta_4^2(n\omega + \frac{\omega}{2})}{\vartheta_3^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[5 \frac{\vartheta_4''(n\omega + \frac{\omega}{2})}{\vartheta_4(n\omega + \frac{\omega}{2})} - 5 \frac{\vartheta_3''(\frac{\omega}{2})}{\vartheta_3(\frac{\omega}{2})} \right. \\ &\quad \left. + 7 \left(\frac{\vartheta'_4(n\omega + \frac{\omega}{2})}{\vartheta_4(n\omega + \frac{\omega}{2})} \right)^2 + 17 \left(\frac{\vartheta'_3(\frac{\omega}{2})}{\vartheta_3(\frac{\omega}{2})} \right)^2 - 24 \frac{\vartheta'_4(n\omega + \frac{\omega}{2})\vartheta'_3(\frac{\omega}{2})}{\vartheta_4(n\omega + \frac{\omega}{2})\vartheta_3(\frac{\omega}{2})} \right], \end{aligned} \quad (13.49)$$

$$\begin{aligned} \frac{d^2}{d\tilde{u}^2} [3(\vartheta_{11}^2 + \vartheta_{12}^2) - \vartheta_{11}\vartheta_{12}] \Big|_{z=\beta} &= 2 \frac{\vartheta_3^2(0)\vartheta_3^2(n\omega + \frac{\omega}{2})}{\vartheta_4^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[5 \frac{\vartheta_3''(n\omega + \frac{\omega}{2})}{\vartheta_3(n\omega + \frac{\omega}{2})} - 5 \frac{\vartheta_4''(\frac{\omega}{2})}{\vartheta_4(\frac{\omega}{2})} \right. \\ &\quad \left. + 7 \left(\frac{\vartheta'_3(n\omega + \frac{\omega}{2})}{\vartheta_3(n\omega + \frac{\omega}{2})} \right)^2 + 17 \left(\frac{\vartheta'_4(\frac{\omega}{2})}{\vartheta_4(\frac{\omega}{2})} \right)^2 - 24 \frac{\vartheta'_3(n\omega + \frac{\omega}{2})\vartheta'_4(\frac{\omega}{2})}{\vartheta_3(n\omega + \frac{\omega}{2})\vartheta_4(\frac{\omega}{2})} \right], \end{aligned} \quad (13.50)$$

$$\begin{aligned} \frac{d^2}{d\tilde{u}^2} [3J(z) - \vartheta_{11}\vartheta_{12}] \Big|_{z=\alpha'} &= 2 \frac{\vartheta_3^2(0)\vartheta_1^2(n\omega + \frac{\omega}{2})}{\vartheta_2^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[5 \frac{\vartheta_1''(n\omega + \frac{\omega}{2})}{\vartheta_1(n\omega + \frac{\omega}{2})} - 5 \frac{\vartheta_2''(\frac{\omega}{2})}{\vartheta_2(\frac{\omega}{2})} \right. \\ &\quad \left. + 7 \left(\frac{\vartheta'_1(n\omega + \frac{\omega}{2})}{\vartheta_1(n\omega + \frac{\omega}{2})} \right)^2 + 17 \left(\frac{\vartheta'_2(\frac{\omega}{2})}{\vartheta_2(\frac{\omega}{2})} \right)^2 - 24 \frac{\vartheta'_1(n\omega + \frac{\omega}{2})\vartheta'_2(\frac{\omega}{2})}{\vartheta_1(n\omega + \frac{\omega}{2})\vartheta_2(\frac{\omega}{2})} \right], \end{aligned} \quad (13.51)$$

$$\begin{aligned}
\frac{d^2}{d\tilde{u}^2}[3J(z) - \vartheta_{11}\vartheta_{12}] \Big|_{z=\beta'} &= 2 \frac{\vartheta_3^2(0)\vartheta_2^2(n\omega + \frac{\omega}{2})}{\vartheta_1^2(\frac{\omega}{2})\vartheta_4^2(n\omega)} \left[5 \frac{\vartheta_2''(n\omega + \frac{\omega}{2})}{\vartheta_2(n\omega + \frac{\omega}{2})} - 5 \frac{\vartheta_1''(\frac{\omega}{2})}{\vartheta_1(\frac{\omega}{2})} \right. \\
&\quad \left. + 7 \left(\frac{\vartheta_2'(n\omega + \frac{\omega}{2})}{\vartheta_2(n\omega + \frac{\omega}{2})} \right)^2 + 17 \left(\frac{\vartheta_1'(\frac{\omega}{2})}{\vartheta_1(\frac{\omega}{2})} \right)^2 - 24 \frac{\vartheta_2'(n\omega + \frac{\omega}{2})\vartheta_1'(\frac{\omega}{2})}{\vartheta_2(n\omega + \frac{\omega}{2})\vartheta_1(\frac{\omega}{2})} \right].
\end{aligned} \tag{13.52}$$

These formulae, combined with (13.13), prove Lemma 12.1.

14. LARGE n ASYMPTOTIC FORMULA FOR h_n

We evaluate the large n asymptotic behavior of h_{nn} and then we use formula (1.29). By (4.7), $h_{nn} = [\mathbf{P}_1]_{12}$, and by (5.18),

$$[\mathbf{P}_1]_{12} = [\mathbf{R}_1]_{12} \left(-\frac{n\pi i}{\gamma} \right), \tag{14.1}$$

hence

$$h_{nn} = [\mathbf{R}_1]_{12} \left(-\frac{n\pi i}{\gamma} \right). \tag{14.2}$$

Furthermore, from (6.5) we obtain that

$$h_{nn} = e^{nl} [\mathbf{T}_1]_{12} \left(-\frac{n\pi i}{\gamma} \right), \tag{14.3}$$

and from (7.2), that

$$h_{nn} = e^{nl} [\mathbf{S}_1]_{12} \left(-\frac{n\pi i}{\gamma} \right). \tag{14.4}$$

It follows from (11.1) that

$$\mathbf{S}_1 = \mathbf{M}_1 + \mathbf{X}_1. \tag{14.5}$$

By (8.33),

$$[\mathbf{M}_1]_{12} = \frac{iA\vartheta_4((n+1)\omega)}{\vartheta_4(n\omega)}, \quad \omega = \frac{\pi(1+\zeta)}{2}, \quad A = \frac{\pi\vartheta_1'(0)}{2\vartheta_1(\omega)}, \tag{14.6}$$

and by (12.9),

$$[\mathbf{X}_1]_{12} = \frac{c(n)}{n} + O(n^{-2}), \tag{14.7}$$

where

$$c(n) = X_\alpha + X_{\alpha'} + X_{\beta'} + X_\beta \tag{14.8}$$

is an explicit quasi-periodic function of n . Therefore,

$$h_{nn} = e^{nl} \left[\frac{iA\vartheta_4((n+1)\omega)}{\vartheta_4(n\omega)} + \frac{c(n)}{n} + O(n^{-2}) \right] \left(-\frac{n\pi i}{\gamma} \right). \tag{14.9}$$

By (2.38),

$$e^{\frac{l}{2}} = \frac{\pi\vartheta_1'(0)}{2e\vartheta_1(\omega)} = \frac{A}{e}, \tag{14.10}$$

hence

$$\begin{aligned} h_{nn} &= \left(\frac{A}{e}\right)^{2n} \left[iA \frac{\vartheta_4((n+1)\omega)}{\vartheta_4(n\omega)} + \frac{c(n)}{n} + O(n^{-2}) \right] \left(-\frac{n\pi i}{\gamma} \right) \\ &= \frac{n\pi A^{2n+1} \vartheta_4((n+1)\omega)}{\gamma e^{2n} \vartheta_4(n\omega)} \left(1 + \frac{c_1(n)}{n} + O(n^{-2}) \right), \end{aligned} \quad (14.11)$$

where

$$c_1(n) = \frac{c(n)\vartheta_4(n\omega)}{iA\vartheta_4((n+1)\omega)}. \quad (14.12)$$

From (1.29) and the Stirling formula we obtain that

$$\frac{h_n}{(n!)^2} = \frac{n^{2n} h_{nn}}{(n!)^2 (2\gamma)^{2n}} = \left(\frac{e}{2\gamma}\right)^{2n} \frac{h_{nn}}{2\pi n} \left(1 - \frac{1}{6n} + O(n^{-2}) \right), \quad (14.13)$$

hence by (14.11),

$$\begin{aligned} \frac{h_n}{(n!)^2} &= \left(\frac{e}{2\gamma}\right)^{2n} \frac{1}{2\pi n} \frac{n\pi A^{2n+1} \vartheta_4((n+1)\omega)}{\gamma e^{2n} \vartheta_4(n\omega)} \left(1 + \frac{c_1(n)}{n} - \frac{1}{6n} + O(n^{-2}) \right) \\ &= G^{2n+1} \frac{\vartheta_4((n+1)\omega)}{\vartheta_4(n\omega)} \left(1 + \frac{c_2(n)}{n} + O(n^{-2}) \right), \end{aligned} \quad (14.14)$$

where

$$G = \frac{A}{2\gamma} = \frac{\pi \vartheta_1'(0)}{4\gamma \vartheta_1(\omega)}, \quad c_2(n) = c_1(n) - \frac{1}{6}. \quad (14.15)$$

Observe that $c_1(n)$ has the form,

$$c_1(n) = f(n\omega, \omega), \quad (14.16)$$

where $f(x, \omega)$ is a real analytic function, periodic with respect to both x and ω , of periods π and 2π , respectively, so that

$$f(x + \pi, \omega) = f(x, \omega), \quad f(x, \omega + 2\pi) = f(x, \omega). \quad (14.17)$$

We can now summarize now the asymptotic formula for $h_n/(n!)^2$.

Proposition 14.1. *As $n \rightarrow \infty$,*

$$\frac{h_n}{(n!)^2} = G^{2n+1} \frac{\vartheta_4((n+1)\omega)}{\vartheta_4(n\omega)} \left[1 + \frac{f_0(n\omega, \omega)}{n} + O(n^{-2}) \right], \quad (14.18)$$

where

$$G = \frac{\pi \vartheta_1'(0)}{4\gamma \vartheta_1(\omega)} \quad (14.19)$$

and

$$f_0(x, \omega) = f(x, \omega) - \frac{1}{6} \quad (14.20)$$

is a real analytic function which satisfies the periodicity conditions

$$f_0(x + \pi, \omega) = f_0(x, \omega), \quad f_0(x, \omega + 2\pi) = f_0(x, \omega). \quad (14.21)$$

By (14.12),

$$f(n\omega, \omega) = \frac{X(n\omega, \omega)\vartheta_4(n\omega)}{iA\vartheta_4((n+1)\omega)}. \quad (14.22)$$

where

$$X \equiv X_\alpha + X_{\alpha'} + X_{\beta'} + X_\beta \quad (14.23)$$

and explicit expressions for $X_\alpha, X_{\alpha'}, X_{\beta'}, X_\beta$ are given in Lemma 12.1.

In the subsequent sections, we carry out a concrete evaluation of $f(n\omega, \omega)$, obtaining that

$$f_0(x, \omega) \equiv 0, \quad (14.24)$$

thus we can improve Proposition 14.1 to the following.

Proposition 14.2. *As $n \rightarrow \infty$,*

$$\frac{h_n}{(n!)^2} = G^{2n+1} \frac{\vartheta_4((n+1)\omega)}{\vartheta_4(n\omega)} \left(1 + O(n^{-2})\right), \quad (14.25)$$

where G is defined in (14.19).

To this end, we will first show that $f(n\omega, \omega)$ does not depend on n .

15. $f(x, \omega)$ IS CONSTANT IN x

Denote

$$z = n\omega + \frac{\omega}{2}, \quad \tilde{f}(x, \omega) \equiv f\left(x - \frac{\omega}{2}, \omega\right), \quad (15.1)$$

so that

$$\tilde{f}(z, \omega) = f(n\omega, \omega). \quad (15.2)$$

To prove that $f(x, \omega)$ is constant in x we will prove the following lemmas.

Lemma 15.1. *The function $\tilde{f}(z, \omega)$ is doubly periodic in z .*

Lemma 15.2. *The function $\tilde{f}(z, \omega)$ is analytic throughout the z -plane.*

From these two lemmas, it follows immediately that $\tilde{f}(z, \omega)$ is constant in z , as it is a doubly periodic entire function of z , and it thus follows that $f(x, \omega)$ is constant in x .

To prove Lemma 15.1, we will check that $\tilde{f}(z + \pi\tau, \omega) = \tilde{f}(z, \omega)$. By (14.12), (14.8),

$$c_1(n) \equiv f(n\omega, \omega) = \tilde{f}(z, \omega) = Y_1 + Y_2 + Y_3 + Y_4, \quad (15.3)$$

where

$$\begin{aligned} Y_1 &= \frac{X_\alpha \vartheta_4(n\omega)}{iA\vartheta_4(n\omega + \omega)} = \frac{\vartheta_1(\omega)\vartheta_3^2(0)\vartheta_4^2(z)}{48\pi\vartheta_1'(0)\vartheta_3^2(\frac{\omega}{2})\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})} \left(C_\alpha + 12\pi\xi_\alpha + \frac{\pi^2\eta_\alpha}{2(\beta' - \alpha)} \right), \\ Y_2 &= \frac{X_{\alpha'} \vartheta_4(n\omega)}{iA\vartheta_4(n\omega + \omega)} = \frac{\vartheta_1(\omega)\vartheta_3^2(0)\vartheta_1^2(z)}{48\pi\vartheta_1'(0)\vartheta_2^2(\frac{\omega}{2})\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})} \left(C_{\alpha'} + 12\pi\xi_{\alpha'} + \frac{\pi^2\eta_{\alpha'}}{2(\beta' - \alpha')} \right), \\ Y_3 &= \frac{X_{\beta'} \vartheta_4(n\omega)}{iA\vartheta_4(n\omega + \omega)} = \frac{\vartheta_1(\omega)\vartheta_3^2(0)\vartheta_2^2(z)}{48\pi\vartheta_1'(0)\vartheta_1^2(\frac{\omega}{2})\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})} \left(C_{\beta'} + 12\pi\xi_{\beta'} + \frac{\pi^2\eta_{\beta'}}{2(\beta' - \alpha)} \right), \\ Y_4 &= \frac{X_\beta \vartheta_4(n\omega)}{iA\vartheta_4(n\omega + \omega)} = \frac{\vartheta_1(\omega)\vartheta_3^2(0)\vartheta_3^2(z)}{48\pi\vartheta_1'(0)\vartheta_4^2(\frac{\omega}{2})\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})} \left(C_\beta + 12\pi\xi_\beta + \frac{\pi^2\eta_\beta}{2(\beta - \alpha')} \right). \end{aligned} \quad (15.4)$$

Observe that Y_1, Y_2, Y_3, Y_4 can be written in the form

$$\begin{aligned} Y_1 &= Q_{11}h_{11}(z) + Q_{12}h_{12}(z) + Q_{13}h_{13}(z) + Q_{14}h_{14}(z), \\ Y_2 &= Q_{21}h_{21}(z) + Q_{22}h_{22}(z) + Q_{23}h_{23}(z) + Q_{24}h_{24}(z), \\ Y_3 &= Q_{31}h_{31}(z) + Q_{32}h_{32}(z) + Q_{33}h_{33}(z) + Q_{34}h_{34}(z), \\ Y_4 &= Q_{41}h_{41}(z) + Q_{42}h_{42}(z) + Q_{43}h_{43}(z) + Q_{44}h_{44}(z), \end{aligned} \quad (15.5)$$

where

$$\begin{aligned} h_{11}(z) &= \frac{\vartheta_4^2(z)}{\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \quad h_{12}(z) = \frac{\vartheta'_4(z)\vartheta_4(z)}{\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \quad h_{13}(z) = \frac{\vartheta'_4(z)^2}{\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \\ h_{14}(z) &= \frac{\vartheta'_4(z)\vartheta_4(z)}{\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \quad h_{21}(z) = \frac{\vartheta_1^2(z)}{\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \quad h_{22}(z) = \frac{\vartheta'_1(z)\vartheta_1(z)}{\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \\ h_{23}(z) &= \frac{\vartheta_1^2(z)}{\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \quad h_{24}(z) = \frac{\vartheta'_1(z)\vartheta_1(z)}{\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \quad h_{31}(z) = \frac{\vartheta_2^2(z)}{\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \\ h_{32}(z) &= \frac{\vartheta'_2(z)\vartheta_2(z)}{\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \quad h_{33}(z) = \frac{\vartheta_2^2(z)}{\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \quad h_{34}(z) = \frac{\vartheta'_2(z)\vartheta_2(z)}{\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \\ h_{41}(z) &= \frac{\vartheta_3^2(z)}{\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \quad h_{42}(z) = \frac{\vartheta'_3(z)\vartheta_3(z)}{\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \quad h_{43}(z) = \frac{\vartheta_3^2(z)}{\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \\ h_{44}(z) &= \frac{\vartheta'_3(z)\vartheta_3(z)}{\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \end{aligned} \quad (15.6)$$

and the numbers Q_{ij} do not depend on z . More specifically,

$$\begin{aligned} Q_{11} &= \frac{\vartheta_1(\omega)\vartheta_3^2(0)}{48\pi\vartheta'_1(0)\vartheta_3^2(\frac{\omega}{2})} \left[C_\alpha + 12\pi \frac{\vartheta'_3(\frac{\omega}{2})}{\vartheta_3(\frac{\omega}{2})} + \frac{\pi^2}{2(\beta' - \alpha)} \left(-5 \frac{\vartheta''_3(\frac{\omega}{2})}{\vartheta_3(\frac{\omega}{2})} + 17 \left(\frac{\vartheta'_3(\frac{\omega}{2})}{\vartheta_3(\frac{\omega}{2})} \right)^2 \right) \right], \\ Q_{12} &= -\frac{\vartheta_1(\omega)\vartheta_3^2(0)}{4\vartheta'_1(0)\vartheta_3^2(\frac{\omega}{2})} - \frac{\pi\vartheta_1(\omega)\vartheta_3^2(0)\vartheta'_3(\frac{\omega}{2})}{4(\beta' - \alpha)\vartheta'_1(0)\vartheta_3^3(\frac{\omega}{2})}, \quad Q_{13} = \frac{7\pi\vartheta_1(\omega)\vartheta_3^2(0)}{96(\beta' - \alpha)\vartheta'_1(0)\vartheta_3^2(\frac{\omega}{2})}, \\ Q_{14} &= \frac{5\pi\vartheta_1(\omega)\vartheta_3^2(0)}{96(\beta' - \alpha)\vartheta'_1(0)\vartheta_3^2(\frac{\omega}{2})}, \end{aligned} \quad (15.7)$$

and similar formulae hold for other Q_{ij} . In particular, notice that

$$\begin{aligned} Q_{13} + Q_{14} &= \frac{\pi\vartheta_1(\omega)\vartheta_3^2(0)}{8(\beta' - \alpha)\vartheta'_1(0)\vartheta_3^2(\frac{\omega}{2})} = \frac{\vartheta_1(\omega)\vartheta_1(\frac{\omega}{2})}{8\vartheta'_1(0)\vartheta_2(\frac{\omega}{2})\vartheta_3(\frac{\omega}{2})\vartheta_4(\frac{\omega}{2})}, \\ Q_{23} + Q_{24} &= -\frac{\pi\vartheta_1(\omega)\vartheta_3^2(0)}{8(\beta - \alpha')\vartheta'_1(0)\vartheta_2^2(\frac{\omega}{2})} = -\frac{\vartheta_1(\omega)\vartheta_4(\frac{\omega}{2})}{8\vartheta'_1(0)\vartheta_1(\frac{\omega}{2})\vartheta_2(\frac{\omega}{2})\vartheta_3(\frac{\omega}{2})}, \\ Q_{33} + Q_{34} &= -\frac{\pi\vartheta_1(\omega)\vartheta_3^2(0)}{8(\beta' - \alpha)\vartheta'_1(0)\vartheta_1^2(\frac{\omega}{2})} = -\frac{\vartheta_1(\omega)\vartheta_3(\frac{\omega}{2})}{8\vartheta'_1(0)\vartheta_1(\frac{\omega}{2})\vartheta_2(\frac{\omega}{2})\vartheta_4(\frac{\omega}{2})}, \\ Q_{43} + Q_{44} &= \frac{\pi\vartheta_1(\omega)\vartheta_3^2(0)}{8(\beta - \alpha')\vartheta'_1(0)\vartheta_4^2(\frac{\omega}{2})} = \frac{\vartheta_1(\omega)\vartheta_2(\frac{\omega}{2})}{8\vartheta'_1(0)\vartheta_1(\frac{\omega}{2})\vartheta_3(\frac{\omega}{2})\vartheta_4(\frac{\omega}{2})}. \end{aligned} \quad (15.8)$$

Observe that all $Q_{ij} = Q_{ij}(\omega)$ are periodic functions of ω of period 2π . It follows from equations (1.39) that the functions

$$h_{j1}(z), \quad j = 1, 2, 3, 4, \quad (15.9)$$

are doubly periodic,

$$h_{j1}(z + \pi) = h_{j1}(z), \quad h_{j1}(z + \pi\tau) = h_{j1}(z), \quad (15.10)$$

while the functions,

$$h_{j2}(z), \quad h_{j3}(z), \quad h_{j4}(z), \quad j = 1, 2, 3, 4 \quad (15.11)$$

satisfy the equations,

$$\begin{aligned} h_{jk}(z + \pi) &= h_{jk}(z), \quad k = 2, 3, 4; \\ h_{j2}(z + \pi\tau) &= h_{j2}(z) - 2ih_{j1}(z), \quad h_{j3}(z + \pi\tau) = h_{j3}(z) - 4ih_{j2}(z) - 4h_{j1}(z), \\ h_{j4}(z + \pi\tau) &= h_{j4}(z) - 4ih_{j3}(z) - 4h_{j1}(z). \end{aligned} \quad (15.12)$$

This implies that the functions $Y_j = Y_j(z)$ for $j = 1, 2, 3, 4$, satisfy the equations

$$\begin{aligned} Y_j(z + \pi) &= Y_j(z), \\ Y_j(z + \pi\tau) &= Y_j(z) + (-2iQ_{j2} - 4Q_{j3} - 4Q_{j4})h_{j1}(z) \\ &\quad + (-4iQ_{j3} - 4iQ_{j4})h_{j2}(z). \end{aligned} \quad (15.13)$$

The proof of Lemma 15.1 then follows immediately from (15.13) and the following identities:

$$(Q_{13} + Q_{14})h_{11}(z) + (Q_{23} + Q_{24})h_{21}(z) + (Q_{33} + Q_{34})h_{31}(z) + (Q_{43} + Q_{44})h_{41}(z) \equiv 0 \quad (15.14)$$

$$(Q_{13} + Q_{14})h_{12}(z) + (Q_{23} + Q_{24})h_{22}(z) + (Q_{33} + Q_{34})h_{32}(z) + (Q_{43} + Q_{44})h_{42}(z) \equiv 0 \quad (15.15)$$

$$Q_{12}h_{11}(z) + Q_{22}h_{21}(z) + Q_{32}h_{31}(z) + Q_{42}h_{41}(z) \equiv 0, \quad (15.16)$$

which are proven below. Introduce here the notation

$$\vartheta_j \equiv \vartheta_j\left(\frac{\omega}{2}\right), \quad \vartheta'_j \equiv \vartheta'_j\left(\frac{\omega}{2}\right), \quad \vartheta''_j \equiv \vartheta''_j\left(\frac{\omega}{2}\right) \quad \text{for } j = 1, 2, 3, 4. \quad (15.17)$$

The sum in (15.14) can be written as

$$\frac{\vartheta_1(\omega) [\vartheta_1^2 \vartheta_4^2(z) - \vartheta_4^2 \vartheta_1^2(z) + \vartheta_2^2 \vartheta_3^2(z) - \vartheta_3^2 \vartheta_2^2(z)]}{8\vartheta'_1(0)\vartheta_1\vartheta_2\vartheta_3\vartheta_4\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}, \quad (15.18)$$

which is zero by the Jacobi identity (E.9).

The sum in (15.15) can be written as

$$\frac{\vartheta_1(\omega) [\vartheta_1^2 \vartheta'_4(z)\vartheta_4(z) - \vartheta_4^2 \vartheta'_1(z)\vartheta_1(z) + \vartheta_2^2 \vartheta'_3(z)\vartheta_3(z) - \vartheta_3^2 \vartheta'_2(z)\vartheta_2(z)]}{8\vartheta'_1(0)\vartheta_1\vartheta_2\vartheta_3\vartheta_4\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})}. \quad (15.19)$$

Using the identities (E.2), we can write the expression in brackets in the numerator of (15.19) as

$$\begin{aligned} &\frac{\vartheta'_1(z)}{\vartheta_1(z)} \left[\vartheta_1^2 \vartheta_4^2(z) - \vartheta_4^2 \vartheta_1^2(z) - \vartheta_3^2 \vartheta_2^2(z) + \vartheta_2^2 \vartheta_3^2(z) \right] \\ &\quad + \frac{\vartheta_2(z)\vartheta_3(z)\vartheta_4(z)}{\vartheta_1(z)} \left[\vartheta_3^2 \vartheta_2^2(0) - \vartheta_1^2 \vartheta_4^2(0) - \vartheta_2^2 \vartheta_3^2(0) \right]. \end{aligned} \quad (15.20)$$

The first term in (15.20) vanishes by (E.9) and the second term vanishes by (E.10). Thus (15.15) is proven.

Finally, we can expand the sum in (15.16) and make the substitutions from identities (E.2), to obtain

$$\begin{aligned} & \frac{\vartheta_1(w)}{4\vartheta'_1(0)\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})} \left[\frac{\vartheta'_1(\vartheta_4^2\vartheta_1^2(z) - \vartheta_1^2\vartheta_4^2(z) - \vartheta_2^2\vartheta_3^2(z) + \vartheta_3^2\vartheta_2^2(z))}{\vartheta_1^2\vartheta_2\vartheta_3\vartheta_4} \right. \\ & \left. + \frac{\vartheta_3^2(z)}{\vartheta_4^2} \left(\vartheta_3^2(0) + \frac{\vartheta_2^2\vartheta_4^2(0)}{\vartheta_1^2} \right) + \frac{\vartheta_1^2(z)}{\vartheta_2^2} \left(\vartheta_3^2(0) - \frac{\vartheta_4^2\vartheta_2^2(0)}{\vartheta_1^2} \right) - \frac{\vartheta_3^2(0)}{\vartheta_1^2} \vartheta_2^2(z) \right]. \end{aligned} \quad (15.21)$$

Once again, the first term vanishes by (E.9). If we substitute the identity (E.11) in the second and third terms, (15.21) becomes simply

$$\frac{\vartheta_1(w)}{4\vartheta'_1(0)\vartheta_1\vartheta_4(z - \frac{\omega}{2})\vartheta_4(z + \frac{\omega}{2})} \left[\vartheta_3^2(z)\vartheta_2^2(0) - \vartheta_2^2(z)\vartheta_3^2(0) - \vartheta_1^2(z)\vartheta_4^2(0) \right], \quad (15.22)$$

which is zero by (E.10). This proves (15.16) and thus Lemma 15.1.

We now turn to the proof of Lemma 15.2. Notice that in the fundamental rectangle

$$\mathcal{R} = \{z \in \mathbb{C} : -\frac{\pi}{2} \leq \operatorname{Re} z \leq \frac{\pi}{2}, 0 \leq \operatorname{Im} z \leq \pi\tau\}, \quad (15.23)$$

the function $\tilde{f}(z, \omega)$ has the two simple poles,

$$z_{1,2} = \frac{\pi\tau}{2} \pm \frac{\omega}{2}. \quad (15.24)$$

Because of the double-periodicity of $\tilde{f}$, we must have

$$\operatorname{Res}_{z=z_1} \tilde{f}(z, \omega) = -\operatorname{Res}_{z=z_2} \tilde{f}(z, \omega). \quad (15.25)$$

Denote

$$R_{ij}(\omega) = \operatorname{Res}_{z=z_1} h_{ij}(z, \omega). \quad (15.26)$$

Then we have that

$$\begin{aligned} R_{11} &= \frac{\vartheta_1^2}{\vartheta'_1(0)\vartheta_1(\omega)}, \quad R_{12} = \frac{(\vartheta'_1 - i\vartheta_1)\vartheta_1}{\vartheta'_1(0)\vartheta_1(\omega)}, \quad R_{13} = \frac{(\vartheta'_1 - i\vartheta_1)^2}{\vartheta'_1(0)\vartheta_1(\omega)}, \quad R_{14} = \frac{(\vartheta''_1 - 2i\vartheta'_1 - \vartheta_1)\vartheta_1}{\vartheta'_1(0)\vartheta_1(\omega)}, \\ R_{21} &= \frac{\vartheta_4^2}{\vartheta'_1(0)\vartheta_1(\omega)}, \quad R_{22} = \frac{(\vartheta'_4 - i\vartheta_4)\vartheta_4}{\vartheta'_1(0)\vartheta_1(\omega)}, \quad R_{23} = \frac{(\vartheta'_4 - i\vartheta_4)^2}{\vartheta'_1(0)\vartheta_1(\omega)}, \quad R_{24} = \frac{(\vartheta''_4 - 2i\vartheta'_4 - \vartheta_4)\vartheta_4}{\vartheta'_1(0)\vartheta_1(\omega)}, \\ R_{31} &= \frac{-\vartheta_3^2}{\vartheta'_1(0)\vartheta_1(\omega)}, \quad R_{32} = \frac{-(\vartheta'_3 - i\vartheta_3)\vartheta_3}{\vartheta'_1(0)\vartheta_1(\omega)}, \quad R_{33} = \frac{-(\vartheta'_3 - i\vartheta_3)^2}{\vartheta'_1(0)\vartheta_1(\omega)}, \quad R_{34} = \frac{-(\vartheta''_3 - 2i\vartheta'_3 - \vartheta_3)\vartheta_3}{\vartheta'_1(0)\vartheta_1(\omega)}, \\ R_{41} &= \frac{-\vartheta_2^2}{\vartheta'_1(0)\vartheta_1(\omega)}, \quad R_{42} = \frac{-(\vartheta'_2 - i\vartheta_2)\vartheta_2}{\vartheta'_1(0)\vartheta_1(\omega)}, \quad R_{43} = \frac{-(\vartheta'_2 - i\vartheta_2)^2}{\vartheta'_1(0)\vartheta_1(\omega)}, \quad R_{44} = \frac{-(\vartheta''_2 - 2i\vartheta'_2 - \vartheta_2)\vartheta_2}{\vartheta'_1(0)\vartheta_1(\omega)}, \end{aligned} \quad (15.27)$$

and

$$\operatorname{Res}_{z=z_1} \tilde{f}(z, \omega) = \sum_{j,k=1}^4 Q_{jk} R_{jk}. \quad (15.28)$$

A priori, the sum in (15.28) is quite complicated, so we evaluate first the imaginary part. Multiplying out (15.28) and again making the substitutions from (E.2), we get

$$\operatorname{Im} \operatorname{Res}_{z=z_1} \tilde{f}(z, \omega) = \frac{(\vartheta_2^4 - \vartheta_4^4)(\vartheta_1^2\vartheta_3^2(0) + \vartheta_2^2\vartheta_4^2(0) - \vartheta_4^2\vartheta_2^2(0))}{4\vartheta'_1(0)^2\vartheta_1^2\vartheta_2^2\vartheta_4^2}, \quad (15.29)$$

which is zero by (E.11).

Substituting the identities (E.5), along with (E.2), into the sum (15.28) gives

$$\operatorname{Res}_{z=z_1} \tilde{f}(z, \omega) = \frac{24A + 17B + 12C + 10D + 7E + 5F}{96\vartheta_1'(0)^2\vartheta_1^2\vartheta_2^3\vartheta_3\vartheta_4^3}, \quad (15.30)$$

where

$$\begin{aligned} A &= \vartheta_3^2(\vartheta_2^4 + \vartheta_4^4) \left[\vartheta_2^2\vartheta_4^2\vartheta_2^2(0)\vartheta_4^2(0) - \vartheta_1^2\vartheta_3^2(0) (\vartheta_2^2\vartheta_4^2(0) - \vartheta_4^2\vartheta_2^2(0)) \right], \\ B &= -\vartheta_3^2 (\vartheta_2^8\vartheta_4^4(0) + \vartheta_4^8\vartheta_2^4(0)), \\ C &= \vartheta_1^2\vartheta_2^2\vartheta_4^2 \left(\vartheta_2^4 + \vartheta_4^4 - \vartheta_1^4 - \vartheta_3^4 \right), \\ D &= -\vartheta_2^4\vartheta_3^2\vartheta_4^4\vartheta_3^4(0), \\ E &= -\vartheta_3^2 \left[\vartheta_1^4\vartheta_3^4(0) (\vartheta_2^4 + \vartheta_4^4) + \vartheta_2^4\vartheta_4^4(\vartheta_2^4(0) + \vartheta_4^4(0)) \right], \\ F &= \vartheta_2^2\vartheta_4^2\vartheta_3^2(0) \left[\vartheta_2^2\vartheta_4^2(0) (\vartheta_3^4 - \vartheta_1^4 - \vartheta_4^4 + \vartheta_2^4) + \vartheta_4^2\vartheta_2^2(0) (\vartheta_4^4 - \vartheta_2^4 + \vartheta_3^4 - \vartheta_1^4) \right]. \end{aligned} \quad (15.31)$$

Note that none of these terms involve derivatives of theta functions. We immediately have $C = 0$ by the identity (E.8). We can also use this identity to write

$$F = 2\vartheta_2^2\vartheta_4^2\vartheta_3^2(0) \left[\vartheta_2^2\vartheta_4^2(0) (\vartheta_3^4 - \vartheta_4^4) + \vartheta_4^2\vartheta_2^2(0) (\vartheta_3^4 - \vartheta_2^4) \right], \quad (15.32)$$

and (E.11) to write

$$A = \vartheta_3^2(\vartheta_2^4 + \vartheta_4^4) \left[\vartheta_2^2\vartheta_4^2\vartheta_2^2(0)\vartheta_4^2(0) + \vartheta_1^4\vartheta_3^2(0) \right]. \quad (15.33)$$

We now combine the terms A, B , and E to obtain

$$\begin{aligned} 24A + 17B + 7E &= 24\vartheta_2^2\vartheta_3^2\vartheta_4^2(\vartheta_4^2\vartheta_2^2(0) - \vartheta_2^2\vartheta_4^2(0))(\vartheta_2^2\vartheta_2^2(0) - \vartheta_4^2\vartheta_4^2(0)) \\ &\quad - 21\vartheta_2^4\vartheta_3^2\vartheta_4^4\vartheta_3^4(0) + 7\vartheta_3^6(\vartheta_2^4 + \vartheta_4^4)(\vartheta_2^4(0) + \vartheta_4^4(0)) \\ &\quad - 7\vartheta_3^2(\vartheta_4^8\vartheta_4^4(0) + \vartheta_2^8\vartheta_2^4(0)). \end{aligned} \quad (15.34)$$

By (E.11) and (E.13) we can write this sum as

$$\begin{aligned} 24A + 17B + 7E &= 24\vartheta_1^2\vartheta_2^2\vartheta_3^2\vartheta_4^2\vartheta_3^2(0)(\vartheta_2^2\vartheta_2^2(0) - \vartheta_4^2\vartheta_4^2(0)) - 21\vartheta_2^4\vartheta_3^2\vartheta_4^4\vartheta_3^4(0) \\ &\quad + 7\vartheta_3^6\vartheta_3^4(0)(\vartheta_2^4 + \vartheta_4^4) - 7\vartheta_3^2(\vartheta_4^8\vartheta_4^4(0) + \vartheta_2^8\vartheta_2^4(0)). \end{aligned} \quad (15.35)$$

We now combine all terms and use (E.11) and (E.13) to write all terms solely in terms of ϑ_2, ϑ_4 , and factors which are constant with respect to ω , yielding

$$\begin{aligned} 24A + 17B + 10D + 7E + 5F &= \left(41 \frac{\vartheta_2^2\vartheta_4^2\vartheta_4^2(0)}{\vartheta_3^2(0)} + 24 \frac{\vartheta_4^4\vartheta_4^4(0)}{\vartheta_2^2(0)\vartheta_3^2(0)} + 17 \frac{\vartheta_2^4\vartheta_2^4(0)}{\vartheta_3^2(0)} \right) \\ &\quad \times \left(\vartheta_2^4(0) + \vartheta_4^4(0) - \vartheta_3^4(0) \right), \end{aligned} \quad (15.36)$$

which is zero by (E.13). Lemma 15.2 is thus proven, and it follows that $\tilde{f}(z, \omega)$ is constant in z . To evaluate the constant, we can take $z = 0$.

16. EVALUATION OF $\tilde{f}(0, \omega)$

We will evaluate $\tilde{f}(z, \omega)$ at $z = 0$. Notice that many of the functions $h_{jk}(z)$ vanish at $z = 0$. In fact we have

$$\tilde{f}(0, \omega) = Q_{11}h_{11}(0) + Q_{14}h_{14}(0) + Q_{23}h_{23}(0) + Q_{31}h_{31}(0) + Q_{34}h_{34}(0) + Q_{41}h_{41}(0) + Q_{44}h_{44}(0). \quad (16.1)$$

The identities (E.4) and (E.3) allow us eliminate all derivatives of theta functions from the sum (16.1) except $\vartheta'_2, \vartheta''_2$. Making these substitutions and simplifying, (16.1) becomes

$$\begin{aligned} \tilde{f}(0, \omega) = & \frac{\vartheta_1(\omega)(\vartheta_1^2\vartheta_4^2(0) + \vartheta_2^2\vartheta_3^2(0) - \vartheta_3^2\vartheta_2^2(0))}{96\vartheta'_1(0)} \left(\frac{5\vartheta''_2(0)}{\vartheta_1\vartheta_2\vartheta_3\vartheta_4^3} + \frac{17(\vartheta'_2)^2}{\vartheta_1\vartheta_2^3\vartheta_3\vartheta_4^3} - \frac{5\vartheta''_2}{\vartheta_1\vartheta_2^2\vartheta_3\vartheta_4^3} \right. \\ & \left. + \frac{24\vartheta'_2(\vartheta_3^2\vartheta_2^2(0) + \vartheta_1^2\vartheta_4^2(0))}{\vartheta_1^2\vartheta_2^3\vartheta_3\vartheta_4^2} \right) + \frac{\vartheta_1(\omega)(24A + 17B + 7C + 5D + 3E + 2F)}{96\vartheta'_1(0)\vartheta_1^3\vartheta_2^3\vartheta_3^3\vartheta_4^3\vartheta_2^2(0)} \end{aligned} \quad (16.2)$$

where

$$\begin{aligned} A &= \vartheta_2^2\vartheta_4^2\vartheta_2^2(0)\vartheta_3^2(0)(\vartheta_1^4\vartheta_4^4(0) + \vartheta_3^4\vartheta_2^4(0)), \\ B &= \vartheta_4^2\vartheta_2^2(0)(\vartheta_1^2\vartheta_4^2(0) - \vartheta_3^2\vartheta_2^2(0))((\vartheta_3^2\vartheta_2^2(0) + \vartheta_1^2\vartheta_4^2(0))^2 - \vartheta_1^2\vartheta_3^2\vartheta_2^2(0)\vartheta_4^2(0)), \\ C &= \vartheta_2^4\vartheta_4^2\vartheta_2^2(0)\vartheta_3^4(0)(\vartheta_1^2\vartheta_4^2(0) - \vartheta_3^2\vartheta_2^2(0)), \\ D &= \vartheta_1^2\vartheta_3^2 \left[\vartheta_4^2\vartheta_2^4(0)\vartheta_4^2(0)(\vartheta_3^2\vartheta_2^2(0) - \vartheta_1^2\vartheta_4^2(0)) + \vartheta_3^2(0) \left(\vartheta_2^2(0)(\vartheta_3^4\vartheta_2^4(0) - \vartheta_1^4\vartheta_4^4(0)) \right. \right. \\ & \quad \left. \left. + \vartheta_2^2\vartheta_1^2(0)\vartheta_4^2(0)(\vartheta_1^2(\vartheta_4^2(0) + \vartheta_2^2\vartheta_3^2(0))) \right) \right], \\ E &= \vartheta_1^2\vartheta_2^2\vartheta_3^2\vartheta_2^2(0)\vartheta_3^2(0) \left[\vartheta_4^2(0)(\vartheta_1^2\vartheta_3^2(0) - \vartheta_4^2\vartheta_2^2(0)) - \vartheta_2^2\vartheta_2^4(0) \right], \\ F &= \vartheta_1^2\vartheta_2^2\vartheta_3^2\vartheta_2^2(0) \left[4\vartheta_2^2\vartheta_3^2(0)\vartheta_4^4(0) - \vartheta_2^2(0)(\vartheta_1^2\vartheta_2^2(0)\vartheta_4^2(0) + \vartheta_3^2\vartheta_4^4(0) + \vartheta_3^2\vartheta_3^4(0)) \right]. \end{aligned} \quad (16.3)$$

The identity (E.10) implies that all terms in (16.2) involving derivatives of theta functions in the numerator vanish. Additionally, (E.10)-(E.13) allow us to simplify the numbers B, C, D and E . Namely, we have

$$\begin{aligned} B &= -\vartheta_2^2\vartheta_4^2\vartheta_2^2(0)\vartheta_3^2(0)((\vartheta_3^2\vartheta_2^2(0) + \vartheta_1^2\vartheta_4^2(0))^2 - \vartheta_1^2\vartheta_3^2\vartheta_2^2(0)\vartheta_4^2(0)), \\ C &= -\vartheta_2^6\vartheta_4^2\vartheta_2^2(0)\vartheta_3^6(0), \quad D = \vartheta_1^2\vartheta_2^2\vartheta_3^2\vartheta_2^2(0)\vartheta_3^2(0) \left(\vartheta_1^2\vartheta_3^2(0)\vartheta_4^2(0) + \vartheta_3^2\vartheta_2^2(0)\vartheta_3^2(0) + \vartheta_4^2\vartheta_2^2(0)\vartheta_4^2(0) \right), \\ E &= -\vartheta_1^2\vartheta_2^4\vartheta_3^2\vartheta_2^2(0)\vartheta_3^6(0). \end{aligned} \quad (16.4)$$

Combining these terms gives us

$$\begin{aligned}
24A + 17B + 7C + 5D + 3E + 2F = \\
\vartheta_2^2 \vartheta_2^2(0) \left[-12\vartheta_1^2 \vartheta_3^2 \vartheta_4^2 \vartheta_2^2(0) \vartheta_3^2(0) \vartheta_4^2(0) + 7\vartheta_4^2 \vartheta_3^2(0) (\vartheta_3^4 \vartheta_2^4(0) + \vartheta_1^4 \vartheta_4^4(0) - \vartheta_2^4 \vartheta_3^4(0)) \right. \\
+ 8\vartheta_1^2 \vartheta_2^2 \vartheta_3^2 \vartheta_3^2(0) \vartheta_4^4(0) + 5\vartheta_1^4 \vartheta_3^2 \vartheta_3^2(0) \vartheta_4^2(0) + 3\vartheta_1^2 \vartheta_3^2 \vartheta_3^2(0) (\vartheta_3^2 \vartheta_2^2(0) - \vartheta_2^2 \vartheta_3^2(0)) \\
\left. - 2\vartheta_1^2 \vartheta_3^2 \vartheta_2^2(0) \vartheta_4^2(0) (\vartheta_1^2 \vartheta_2^2(0) + \vartheta_3^2 \vartheta_4^2(0)) \right]. \tag{16.5}
\end{aligned}$$

Again using (E.10)-(E.13), this expression simplifies to

$$24A + 17B + 7C + 5D + 3E + 2F = 8\vartheta_1^2 \vartheta_2^2 \vartheta_3^2 \vartheta_4^2 \vartheta_2^2(0) \vartheta_3^2(0) \vartheta_4^2(0). \tag{16.6}$$

Inserting this into (16.2), we get

$$\tilde{f}(0, \omega) \equiv \tilde{f}(z, \omega) \equiv f(n\omega, \omega) = \frac{\vartheta_1(\omega) \vartheta_2^2(0) \vartheta_3^2(0) \vartheta_4^2(0)}{12\vartheta_1'(0) \vartheta_1 \vartheta_2 \vartheta_3 \vartheta_4} = \frac{1}{6} \tag{16.7}$$

by (E.1) and (E.6). It then follows that

$$f_0(n\omega, \omega) = f(n\omega, \omega) - \frac{1}{6} = 0. \tag{16.8}$$

This proves (14.24) and therefore Proposition 14.2.

17. LARGE n ASYMPTOTICS OF Z_n

By substituting (14.25) into (1.24) we obtain that

$$\begin{aligned}
\frac{\tau_n}{\prod_{k=0}^{n-1} (k!)^2} &= 2^{n^2} \prod_{k=0}^{n-1} \frac{h_k}{(k!)^2} \\
&= 2^{n^2} h_0 \prod_{k=1}^{n-1} \left[G^{2k+1} \frac{\vartheta_4((k+1)\omega)}{\vartheta_4(k\omega)} (1 + O(k^{-2})) \right] \\
&= C \vartheta_4(n\omega) (2G)^{n^2} \left(1 + O(n^{-1}) \right), \tag{17.1}
\end{aligned}$$

where $C > 0$ does not depend on n . Thus, by (1.15),

$$Z_n = \frac{[\sinh(\gamma - t) \sinh(\gamma + t)]^{n^2} \tau_n}{\left(\prod_{k=0}^{n-1} k! \right)^2} = C \vartheta_4(n\omega) F^{n^2} \left(1 + O(n^{-1}) \right), \tag{17.2}$$

where

$$F = 2G \sinh(\gamma - t) \sinh(\gamma + t) = \frac{\pi \sinh(\gamma - t) \sinh(\gamma + t) \vartheta_1'(0)}{2\gamma \vartheta_1(\omega)}. \tag{17.3}$$

Theorem 1.1 is proved.

APPENDIX A. PROOF OF FORMULA (1.20) FOR τ_n

We have

$$\tau_n = \det \left(\phi^{(i+j-2)}(t) \right)_{1 \leq i, j \leq n} \quad (\text{A.1})$$

where

$$\begin{aligned} \phi(t) &= \frac{\sinh(2\gamma)}{\sinh(\gamma+t)\sinh(\gamma-t)} = \frac{2(1-e^{4\gamma})}{(1-e^{-2(\gamma-t)})(1-e^{-2(\gamma+t)})} \\ &= \frac{2}{1-e^{-2(\gamma-t)}} + \frac{2}{1-e^{-2(\gamma+t)}} - 2 = 2 \sum_{l=0}^{\infty} e^{-2l(\gamma-t)} + 2 \sum_{l=0}^{\infty} e^{-2l(\gamma+t)} - 2 \\ &= 2 \sum_{l=0}^{\infty} e^{-2l(\gamma-t)} + 2 \sum_{l=1}^{\infty} e^{-2l(\gamma+t)} = 2 \sum_{l=-\infty}^{\infty} e^{2tl} e^{-2\gamma|l|}. \end{aligned} \quad (\text{A.2})$$

Thus,

$$\phi^{(k)}(t) = 2 \sum_{l=-\infty}^{\infty} (2l)^k e^{2tl} e^{-2\gamma|l|}. \quad (\text{A.3})$$

From equation (A.3) and multi-linearity of the determinant function, we have that τ_n is equal to

$$\begin{aligned} 2^n \sum_{l_1, l_2, \dots, l_n = -\infty}^{\infty} \det \begin{pmatrix} e^{2tl_1-2\gamma|l_1|} & (2l_1)e^{2tl_1-2\gamma|l_1|} & \dots & (2l_1)^{n-1}e^{2tl_1-2\gamma|l_1|} \\ (2l_2)e^{2tl_2-2\gamma|l_2|} & (2l_2)^2e^{2tl_2-2\gamma|l_2|} & \dots & (2l_2)^ne^{2tl_2-2\gamma|l_2|} \\ (2l_3)^2e^{2tl_3-2\gamma|l_3|} & (2l_3)^3e^{2tl_3-2\gamma|l_3|} & \dots & (2l_3)^{n+1}e^{2tl_3-2\gamma|l_3|} \\ \vdots & \vdots & \ddots & \vdots \\ (2l_n)^{n-1}e^{2tl_n-2\gamma|l_n|} & (2l_n)^ne^{2tl_n-2\gamma|l_n|} & \dots & (2l_n)^{2n-2}e^{2tl_n-2\gamma|l_n|} \end{pmatrix} \\ = 2^n \sum_{l_1, l_2, \dots, l_n = -\infty}^{\infty} \Delta(2l) \prod_{i=1}^n e^{2tl_i-2\gamma|l_i|} \prod_{i=1}^n (2l_i)^{i-1} \\ = 2^{n^2} \sum_{l_1, l_2, \dots, l_n = -\infty}^{\infty} \prod_{i=1}^n \Delta(l) e^{2tl_i-2\gamma|l_i|} \prod_{i=1}^n (l_i)^{i-1}, \end{aligned} \quad (\text{A.4})$$

where $\Delta(l) = \prod_{i < j} (l_i - l_j)$ is the Vandermonde determinant. Note that, up to sign, this expression for τ_n is invariant with respect to any permutation of l_i 's. So, multiplying by their signs and then summing over all permutations, we get

$$n! \tau_n = 2^{n^2} \sum_{l_1, l_2, \dots, l_n = -\infty}^{\infty} \prod_{i=1}^n e^{2tl_i-2\gamma|l_i|} \Delta(l_i) \sum_{\pi \in S_n} (-1)^\pi \prod_{i=1}^n (l_i)^{\pi(i)-1}, \quad (\text{A.5})$$

thus

$$\tau_n = \frac{2^{n^2}}{n!} \sum_{l_1, l_2, \dots, l_n = -\infty}^{\infty} \Delta(l_i)^2 \prod_{i=1}^n e^{2tl_i-2\gamma|l_i|}. \quad (\text{A.6})$$

APPENDIX B. DERIVATION OF FORMULA (1.24)

Multilinearity of the determinant function, combined with the form of the Vandermonde matrix, allows us to replace $\Delta(l)$ with

$$\det \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 \\ P_1(l_1) & P_1(l_2) & P_1(l_3) & \cdots & P_1(l_n) \\ P_2(l_1) & P_2(l_2) & P_2(l_3) & \cdots & P_2(l_n) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ P_{n-1}(l_1) & P_{n-1}(l_2) & P_{n-1}(l_3) & \cdots & P_{n-1}(l_n) \end{pmatrix}, \quad (\text{B.1})$$

where $\{P_j(x)\}_{j=0}^\infty$ is the system of monic polynomials orthogonal with respect to the weight $w(l)$. Then (1.20) becomes

$$\tau_n = \frac{2^{n^2}}{n!} \sum_{l_1, \dots, l_n=1}^\infty \left(\sum_{\pi \in S_n} (-1)^\pi \prod_{k=1}^n P_{\pi(k)-1}(l_k) \right)^2 \prod_{k=1}^n w(l_k). \quad (\text{B.2})$$

The orthogonality condition ensures that, after summing, only diagonal terms are non-zero, so we get

$$\tau_n = \frac{2^{n^2}}{n!} \sum_{l_1, \dots, l_n=1}^\infty \left(\sum_{\pi \in S_n} \prod_{k=1}^n P_{\pi(k)-1}^2(l_k) \right) \prod_{k=1}^n w(l_k) = 2^{n^2} \prod_{k=0}^{n-1} h_k. \quad (\text{B.3})$$

APPENDIX C. PROOF OF (10.5)

From, (10.4), (7.1), (7.2), and (2.32), we have that the jump $j_{\tilde{Q}}$ on $(\alpha' - \varepsilon, \alpha')$ is given by

$$\begin{aligned} j_{\tilde{Q}} &= e^{\frac{in\pi z}{2\gamma}\sigma_3} e^{-n(g_-(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} j_S e^{n(g_+(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} e^{\frac{in\pi z}{2\gamma}\sigma_3} \\ &= e^{\frac{in\pi z}{2\gamma}\sigma_3} e^{-n(g_-(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} j_-^{-1} j_T j_+^{-1} e^{n(g_+(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} e^{\frac{in\pi z}{2\gamma}\sigma_3} \\ &= e^{\frac{in\pi z}{2\gamma}\sigma_3} e^{-n(g_-(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} e^{n(g_+(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} e^{\frac{in\pi z}{2\gamma}\sigma_3} \\ &= e^{\frac{in\pi z}{2\gamma}\sigma_3} \begin{pmatrix} 0 & e^{-n(g_+(z) + g_-(z) - V(z) - l)} \\ -e^{n(g_+(z) + g_-(z) - V(z) - l)} & 0 \end{pmatrix} e^{\frac{in\pi z}{2\gamma}\sigma_3} \\ &= \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \end{aligned} \quad (\text{C.1})$$

From, (10.4), (7.2), (6.2), and (5.21), we have that the jump $j_{\tilde{Q}}$ on $(\alpha', \alpha' + \varepsilon)$ is given by

$$\begin{aligned}
j_{\tilde{Q}} &= e^{\frac{in\pi z}{2\gamma}\sigma_3} e^{-n(g_-(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} j_S e^{n(g_+(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} e^{\frac{in\pi z}{2\gamma}\sigma_3} \\
&= e^{\frac{in\pi z}{2\gamma}\sigma_3} e^{-n(g_-(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} \begin{pmatrix} \frac{\gamma}{n\pi i} e^{\frac{in\pi z}{2\gamma}} & 0 \\ 0 & \frac{n\pi i}{\gamma} e^{-\frac{in\pi z}{2\gamma}} \end{pmatrix}^{-1} j_T \begin{pmatrix} -\frac{\gamma}{n\pi i} e^{-\frac{in\pi z}{2\gamma}} & 0 \\ 0 & -\frac{n\pi i}{\gamma} e^{\frac{in\pi z}{2\gamma}} \end{pmatrix} \\
&\quad \times e^{n(g_+(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} e^{\frac{in\pi z}{2\gamma}\sigma_3} \\
&= e^{-n(g_-(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} \begin{pmatrix} \frac{n\pi i}{\gamma} & 0 \\ 0 & \frac{\gamma}{n\pi i} \end{pmatrix} e^{n(g_-(z) - \frac{l}{2})\sigma_3} j_R e^{-n(g_+(z) - \frac{l}{2})\sigma_3} \begin{pmatrix} -\frac{\gamma}{n\pi i} & 0 \\ 0 & -\frac{n\pi i}{\gamma} \end{pmatrix} \\
&\quad \times e^{n(g_+(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} \\
&= e^{n\frac{V(z)}{2}\sigma_3} \begin{pmatrix} \frac{n\pi i}{\gamma} & 0 \\ 0 & \frac{\gamma}{n\pi i} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ (\frac{n\pi i}{\gamma})^2 e^{nV(z)} & 1 \end{pmatrix} \begin{pmatrix} -\frac{\gamma}{n\pi i} & 0 \\ 0 & -\frac{n\pi i}{\gamma} \end{pmatrix} e^{-n\frac{V(z)}{2}\sigma_3} \\
&= e^{n\frac{V(z)}{2}\sigma_3} \begin{pmatrix} -1 & 0 \\ -e^{nV(z)} & -1 \end{pmatrix} e^{-n\frac{V(z)}{2}\sigma_3} = \begin{pmatrix} -1 & 0 \\ -1 & -1 \end{pmatrix}.
\end{aligned} \tag{C.2}$$

From (10.4), (7.2), (6.2), (5.21), and analytic continuation of (2.32) into a neighborhood of $[\alpha, \alpha']$, we have that the jump $j_{\tilde{Q}}$ on $(\alpha', \alpha' + i\varepsilon)$ is given by

$$\begin{aligned}
j_{\tilde{Q}} &= e^{-n(g_+(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} \begin{pmatrix} -\frac{n\pi i}{\gamma} & 0 \\ 0 & -\frac{\gamma}{n\pi i} \end{pmatrix} e^{n(g_+(z) - \frac{l}{2})\sigma_3} j_R e^{-n(g_+(z) - \frac{l}{2})\sigma_3} j_+(z)^{-1} \\
&\quad \times e^{n(g_+(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} e^{\frac{in\pi z}{2\gamma}\sigma_3} \\
&= e^{n\frac{V(z)}{2}\sigma_3} \begin{pmatrix} -\frac{n\pi i}{\gamma} & 0 \\ 0 & -\frac{\gamma}{n\pi i} \end{pmatrix} j_R e^{-n(g_+(z) - \frac{l}{2})\sigma_3} j_+(z)^{-1} e^{n(g_+(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} e^{\frac{in\pi z}{2\gamma}\sigma_3} \\
&= e^{n\frac{V(z)}{2}\sigma_3} \begin{pmatrix} -\frac{n\pi i}{\gamma} & 0 \\ 0 & -\frac{\gamma}{n\pi i} \end{pmatrix} \begin{pmatrix} \Pi(z) & \frac{\gamma}{n\pi i} e^{-nV(z)} e^{\frac{in\pi z}{2\gamma}} \\ -\frac{n\pi i}{\gamma} e^{nV(z)} e^{\frac{in\pi z}{2\gamma}} & -\frac{n\pi i}{\gamma} e^{\frac{in\pi z}{2\gamma}} \end{pmatrix} \\
&\quad \times e^{-n(g_+(z) - \frac{l}{2})\sigma_3} \begin{pmatrix} 1 & 0 \\ -e^{-n(g_+(z) - g_-(z))} & 1 \end{pmatrix} e^{n(g_+(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} e^{\frac{in\pi z}{2\gamma}\sigma_3} \\
&= e^{n\frac{V(z)}{2}\sigma_3} \begin{pmatrix} -2i \sin(\frac{n\pi z}{2\gamma}) & -e^{-nV(z)} e^{\frac{in\pi z}{2\gamma}} \\ e^{nV(z)} e^{\frac{in\pi z}{2\gamma}} & e^{\frac{in\pi z}{2\gamma}} \end{pmatrix} \begin{pmatrix} e^{-n\frac{V(z)}{2}} & 0 \\ -e^{n(g_+(z) + g_-(z) - l - \frac{V(z)}{2})} & e^{n\frac{V(z)}{2}} \end{pmatrix} e^{\frac{in\pi z}{2\gamma}\sigma_3} \\
&= \begin{pmatrix} -2i \sin(\frac{n\pi z}{2\gamma}) e^{n\frac{V(z)}{2}} & -e^{-n\frac{V(z)}{2}} e^{\frac{in\pi z}{2\gamma}} \\ e^{n\frac{V(z)}{2}} e^{\frac{in\pi z}{2\gamma}} & e^{-n\frac{V(z)}{2}} e^{\frac{in\pi z}{2\gamma}} \end{pmatrix} \begin{pmatrix} e^{-n\frac{V(z)}{2}} & 0 \\ -e^{n(g_+(z) + g_-(z) - l - \frac{V(z)}{2})} & e^{n\frac{V(z)}{2}} \end{pmatrix} e^{\frac{in\pi z}{2\gamma}\sigma_3} \\
&= \begin{pmatrix} -2i \sin(\frac{n\pi z}{2\gamma}) + e^{\frac{in\pi z}{2\gamma}} e^{n(g_+(z) + g_-(z) - l - V(z))} & -e^{\frac{in\pi z}{2\gamma}} \\ e^{\frac{in\pi z}{2\gamma}} - e^{\frac{in\pi z}{2\gamma}} e^{n(g_+(z) + g_-(z) - l - V(z))} & e^{\frac{in\pi z}{2\gamma}} \end{pmatrix} e^{\frac{in\pi z}{2\gamma}\sigma_3} \\
&= \begin{pmatrix} e^{-\frac{in\pi z}{2\gamma}} & -e^{\frac{in\pi z}{2\gamma}} \\ 0 & e^{\frac{in\pi z}{2\gamma}} \end{pmatrix} e^{\frac{in\pi z}{2\gamma}\sigma_3} = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}.
\end{aligned} \tag{C.3}$$

Similarly, we have that the jump $j_{\tilde{Q}}$ on $(\alpha' - i\varepsilon, \alpha')$ is given by

$$\begin{aligned}
j_{\tilde{Q}} &= e^{\frac{in\pi z}{2\gamma}\sigma_3} e^{-n(g_-(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} j_S e^{n(g_-(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} e^{-\frac{in\pi z}{2\gamma}\sigma_3} \\
&= e^{\frac{in\pi z}{2\gamma}\sigma_3} e^{-n(g_-(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} \begin{pmatrix} \frac{\gamma}{n\pi i} e^{\frac{in\pi z}{2\gamma}} & 0 \\ 0 & \frac{n\pi i}{\gamma} e^{-\frac{in\pi z}{2\gamma}} \end{pmatrix}^{-1} j_T j_-(z) e^{n(g_-(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} e^{-\frac{in\pi z}{2\gamma}\sigma_3} \\
&= e^{n\frac{V(z)}{2}\sigma_3} \begin{pmatrix} \frac{n\pi i}{\gamma} & 0 \\ 0 & \frac{\gamma}{n\pi i} \end{pmatrix} j_R e^{-n(g_-(z) - \frac{l}{2})\sigma_3} j_-(z) e^{n(g_-(z) - \frac{V(z)}{2} - \frac{l}{2})\sigma_3} e^{-\frac{in\pi z}{2\gamma}\sigma_3} \\
&= e^{n\frac{V(z)}{2}\sigma_3} \begin{pmatrix} \frac{n\pi i}{\gamma} & 0 \\ 0 & \frac{\gamma}{n\pi i} \end{pmatrix} \begin{pmatrix} \Pi(z) & \frac{\gamma}{n\pi i} e^{-nV(z)} e^{-\frac{in\pi z}{2\gamma}} \\ -\frac{n\pi i}{\gamma} e^{nV(z)} e^{-\frac{in\pi z}{2\gamma}} & \frac{n\pi i}{\gamma} e^{-\frac{in\pi z}{2\gamma}} \end{pmatrix} \\
&\quad \times \begin{pmatrix} e^{-n\frac{V(z)}{2}} & 0 \\ e^{n(g_+(z) + g_-(z) - l - \frac{V(z)}{2})} & e^{n\frac{V(z)}{2}} \end{pmatrix} e^{-\frac{in\pi z}{2\gamma}\sigma_3} \\
&= \begin{pmatrix} 2i \sin(\frac{n\pi z}{2\gamma}) e^{n\frac{V(z)}{2}} & e^{-n\frac{V(z)}{2}} e^{-\frac{in\pi z}{2\gamma}} \\ -e^{n\frac{V(z)}{2}} e^{-\frac{in\pi z}{2\gamma}} & e^{-n\frac{V(z)}{2}} e^{-\frac{in\pi z}{2\gamma}} \end{pmatrix} \begin{pmatrix} e^{-n\frac{V(z)}{2}} & 0 \\ e^{n(g_+(z) + g_-(z) - l - \frac{V(z)}{2})} & e^{n\frac{V(z)}{2}} \end{pmatrix} e^{-\frac{in\pi z}{2\gamma}\sigma_3} \\
&= \begin{pmatrix} 2i \sin(\frac{n\pi z}{2\gamma}) + e^{-\frac{in\pi z}{2\gamma}} e^{n(g_+(z) + g_-(z) - l - V(z))} & e^{-\frac{in\pi z}{2\gamma}} \\ -e^{-\frac{in\pi z}{2\gamma}} + e^{-\frac{in\pi z}{2\gamma}} e^{n(g_+(z) + g_-(z) - l - V(z))} & e^{-\frac{in\pi z}{2\gamma}} \end{pmatrix} e^{-\frac{in\pi z}{2\gamma}\sigma_3} \\
&= \begin{pmatrix} e^{\frac{in\pi z}{2\gamma}} & e^{-\frac{in\pi z}{2\gamma}} \\ 0 & e^{-\frac{in\pi z}{2\gamma}} \end{pmatrix} e^{-\frac{in\pi z}{2\gamma}\sigma_3} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.
\end{aligned} \tag{C.4}$$

APPENDIX D. PROOF OF PROPOSITION 8.1

From (3.16) and (8.7), we have

$$\tilde{u}_\infty = \frac{\pi}{4}(1 - \zeta) \tag{D.1}$$

This, combined with formula (7.5) for Ω_n immediately gives

$$\frac{\vartheta_3(\tilde{u}_\infty + d)\vartheta_3(-\tilde{u}_\infty + d + \frac{\Omega_n}{2})}{\vartheta_3(-\tilde{u}_\infty + d)\vartheta_3(\tilde{u}_\infty + d + \frac{\Omega_n}{2})} = \frac{\vartheta_3(0)\vartheta_4((n+1)\omega)}{\vartheta_3(\omega)\vartheta_4(n\omega)}. \tag{D.2}$$

Formulae (2.16) give that

$$\frac{(\beta - \beta') + (\alpha' - \alpha)}{4i} = \frac{\pi\vartheta_4^2(0)}{4i} \left[\frac{\vartheta_1^2(\frac{\omega}{2})\vartheta_4^2(\frac{\omega}{2}) + \vartheta_2^2(\frac{\omega}{2})\vartheta_3^2(\frac{\omega}{2})}{\vartheta_1(\frac{\omega}{2})\vartheta_2(\frac{\omega}{2})\vartheta_3(\frac{\omega}{2})\vartheta_4(\frac{\omega}{2})} \right]. \tag{D.3}$$

Plugging the duplication formulae (E.6) and (E.7) into (D.3) yields

$$\frac{(\beta - \beta') + (\alpha' - \alpha)}{4i} = \frac{\pi}{2i} \frac{\vartheta_2(0)\vartheta_4(0)\vartheta_3(\omega)}{\vartheta_1(\omega)}. \tag{D.4}$$

Combining (D.2) and (D.4), we can write the [12] entry of (8.32) as

$$\begin{aligned} [\mathbf{M}_1]_{12} &= \frac{i\pi}{2} \frac{\vartheta_4((n+1)\omega)}{\vartheta_4(n\omega)} \frac{\vartheta_3(0)}{\vartheta_3(\omega)} \frac{\vartheta_2(0)\vartheta_4(0)\vartheta_3(\omega)}{\vartheta_1(\omega)} \\ &= \frac{i\pi}{2} \frac{\vartheta_4((n+1)\omega)}{\vartheta_4(n\omega)} \frac{\vartheta'_1(0)}{\vartheta_2(\frac{\pi\zeta}{2})} = \frac{iA\vartheta_4((n+1)\omega)}{\vartheta_4(n\omega)}. \end{aligned} \quad (\text{D.5})$$

Similarly, we can write the [21] entry of (8.32) as

$$[\mathbf{M}_1]_{21} = \frac{A\vartheta_4(n\omega)}{i\vartheta_4((n-1)\omega)}. \quad (\text{D.6})$$

APPENDIX E. THETA FUNCTION IDENTITIES

The following identities (see [30]) are used in this paper. There are the identities involving derivatives of theta functions:

$$\vartheta'_1(0) = \vartheta_2(0)\vartheta_3(0)\vartheta_4(0), \quad (\text{E.1})$$

$$\begin{aligned} \vartheta'_4(z) &= \frac{\vartheta'_1(z)\vartheta_4(z) - \vartheta_4(0)^2\vartheta_2(z)\vartheta_3(z)}{\vartheta_1(z)}, \\ \vartheta'_2(z) &= \frac{\vartheta'_1(z)\vartheta_2(z) - \vartheta_2(0)^3\vartheta_3(z)\vartheta_4(z)}{\vartheta_1(z)}, \\ \vartheta'_3(z) &= \frac{\vartheta'_1(z)\vartheta_3(z) - \vartheta_3(0)^2\vartheta_2(z)\vartheta_4(z)}{\vartheta_1(z)}, \end{aligned} \quad (\text{E.2})$$

$$\begin{aligned} \vartheta'_4(z) &= \frac{\vartheta'_2(z)\vartheta_4(z) + \vartheta_1(z)\vartheta_3(z)\vartheta_3^2(0)}{\vartheta_2(z)}, \\ \vartheta'_1(z) &= \frac{\vartheta'_2(z)\vartheta_1(z) + \vartheta_3(z)\vartheta_4(z)\vartheta_2^2(0)}{\vartheta_2(z)}, \\ \vartheta'_3(z) &= \frac{\vartheta'_2(z)\vartheta_3(z) + \vartheta_1(z)\vartheta_4(z)\vartheta_4^2(0)}{\vartheta_2(z)}, \end{aligned} \quad (\text{E.3})$$

$$\begin{aligned} \vartheta''_4(z) &= \frac{\vartheta''_2(z)\vartheta_4(z)}{\vartheta_2(z)} + \frac{2\vartheta'_2(z)\vartheta_1(z)\vartheta_3(z)\vartheta_3^2(0)}{\vartheta_2^2(z)} + \frac{\vartheta_4(z)\vartheta_3^2(0)}{\vartheta_2^2(z)} \left(\vartheta_3^2(z)\vartheta_2^2(0) + \vartheta_1^2(z)\vartheta_4^2(0) \right), \\ \vartheta''_1(z) &= \frac{\vartheta''_2(z)\vartheta_1(z)}{\vartheta_2(z)} + \frac{2\vartheta'_2(z)\vartheta_3(z)\vartheta_4(z)\vartheta_2^2(0)}{\vartheta_2^2(z)} + \frac{\vartheta_1(z)\vartheta_2^2(0)}{\vartheta_2^2(z)} \left(\vartheta_3^2(z)\vartheta_3^2(0) + \vartheta_4^2(z)\vartheta_4^2(0) \right), \\ \vartheta''_3(z) &= \frac{\vartheta''_2(z)\vartheta_3(z)}{\vartheta_2(z)} + \frac{2\vartheta'_2(z)\vartheta_1(z)\vartheta_4(z)\vartheta_4^2(0)}{\vartheta_2^2(z)} + \frac{\vartheta_3(z)\vartheta_3^2(0)}{\vartheta_2^2(z)} \left(\vartheta_4^2(z)\vartheta_2^2(0) + \vartheta_1^2(z)\vartheta_4^2(0) \right), \end{aligned} \quad (\text{E.4})$$

$$\begin{aligned}
\vartheta_4''(z) &= \frac{\vartheta_1''(z)\vartheta_4(z)}{\vartheta_1(z)} - \frac{2\vartheta_1'(z)\vartheta_2(z)\vartheta_3(z)\vartheta_4^2(0)}{\vartheta_1^2(z)} + \frac{\vartheta_4^2(0)\vartheta_4^2(z)}{\vartheta_1^2(z)} \left(\vartheta_3^2(z)\vartheta_2^2(0) + \vartheta_2^2(z)\vartheta_3^2(0) \right), \\
\vartheta_2''(z) &= \frac{\vartheta_1''(z)\vartheta_2(z)}{\vartheta_1(z)} - \frac{2\vartheta_1'(z)\vartheta_3(z)\vartheta_4(z)\vartheta_2^2(0)}{\vartheta_1^2(z)} + \frac{\vartheta_2^2(0)\vartheta_2^2(z)}{\vartheta_1^2(z)} \left(\vartheta_4^2(z)\vartheta_3^2(0) + \vartheta_3^2(z)\vartheta_4^2(0) \right), \\
\vartheta_3''(z) &= \frac{\vartheta_1''(z)\vartheta_3(z)}{\vartheta_1(z)} - \frac{2\vartheta_1'(z)\vartheta_2(z)\vartheta_4(z)\vartheta_3^2(0)}{\vartheta_1^2(z)} + \frac{\vartheta_3^2(0)\vartheta_3^2(z)}{\vartheta_1^2(z)} \left(\vartheta_4^2(z)\vartheta_2^2(0) + \vartheta_2^2(z)\vartheta_4^2(0) \right),
\end{aligned} \tag{E.5}$$

the duplication formulae

$$\vartheta_1(2z) = \frac{2\vartheta_1(z)\vartheta_2(z)\vartheta_3(z)\vartheta_4(z)}{\vartheta_1'(0)}, \tag{E.6}$$

$$\vartheta_3(2z)\vartheta_3(0)\vartheta_2^2(0) = \vartheta_1(z)^2\vartheta_4(z)^2 + \vartheta_2(z)^2\vartheta_3(z)^2 \tag{E.7}$$

$$\vartheta_4(2z)\vartheta_4^3(0) = \vartheta_3^4(z) - \vartheta_2^4(z) = \vartheta_4^4(z) - \vartheta_1^4(z), \tag{E.8}$$

the addition formula

$$\begin{aligned}
\vartheta_3(y+z)\vartheta_3(y-z)\vartheta_2^2(0) &= \vartheta_3^2(y)\vartheta_2^2(z) + \vartheta_4^2(y)\vartheta_1^2(z) \\
&= \vartheta_1^2(y)\vartheta_4^2(z) + \vartheta_2^2(y)\vartheta_3^2(z),
\end{aligned} \tag{E.9}$$

and the identities relating squares of theta functions

$$\vartheta_1^2(z)\vartheta_4^2(0) = \vartheta_3^2(z)\vartheta_2^2(0) - \vartheta_2^2(z)\vartheta_3^2(0), \tag{E.10}$$

$$\vartheta_2^2(z)\vartheta_4^2(0) = \vartheta_4^2(z)\vartheta_2^2(0) - \vartheta_1^2(z)\vartheta_3^2(0), \tag{E.11}$$

$$\vartheta_3^2(z)\vartheta_4^2(0) = \vartheta_4^2(z)\vartheta_3^2(0) - \vartheta_1^2(z)\vartheta_2^2(0), \tag{E.12}$$

$$\vartheta_4^2(z)\vartheta_4^2(0) = \vartheta_3^2(z)\vartheta_3^2(0) - \vartheta_2^2(z)\vartheta_2^2(0). \tag{E.13}$$

REFERENCES

- [1] D. Allison and N. Reshetikhin, Numerical study of the 6-vertex model with domain wall boundary conditions, *Ann. Inst. Fourier (Grenoble)* **55** (2005) 1847–1869.
- [2] R. Baxter, *Exactly solved models in statistical mechanics*, Academic Press, San Diego, CA.
- [3] M.T. Batchelor, R.J. Baxter, M.J. O’Rourke, and C.M. Yung, Exact solution and interfacial tension of the six-vertex model with antiperiodic boundary conditions. *J. Phys. A* **28** (1995) 2759–2770.
- [4] P.M. Bleher and V.V. Fokin, Exact solution of the six-vertex model with domain wall boundary conditions. Disordered phase. *Commun. Math. Phys.* **268** (2006), 223–284.
- [5] J. Baik, T. Kriecherbauer, K. T.-R. McLaughlin, and P.D. Miller, Discrete orthogonal polynomials. Asymptotics and applications. *Ann. Math. Studies* **164**. Princeton University Press. Princeton and Oxford, 2007.
- [6] H.J. Brascamp, H. Kunz, and F.Y. Wu, Some rigorous results for the vertex model in statistical mechanics, *J. Math. Phys.* **14** (1973), 1927–1932.
- [7] P.M. Bleher and K. Liechty, Exact solution of the six-vertex model with domain wall boundary conditions. Ferroelectric phase, *Commun. Math. Phys.* **286** (2009), 777–801.
- [8] P.M. Bleher and K. Liechty, Exact solution of the six-vertex model with domain wall boundary condition. Critical line between ferroelectric and disordered phases, *J. Statist. Phys.* **134** (2009), 463–485.
- [9] A. Borodin and D. Boyarchenko, Distribution of the first particle in discrete orthogonal polynomial ensembles, *Commun. Math. Phys.* **234** (2003), 287–338.
- [10] D.M. Bressoud, *Proofs and Confirmations: the Story of the Alternating Sign Matrix Conjecture*, published jointly by the Mathematical Association of America (Spectrum Series) and Cambridge University Press (or Cambridge University Press, NY) 1999.

- [11] F. Colomo and A.G. Pronko, Square ice, alternating sign matrices, and classical orthogonal polynomials, *J. Stat. Mech. Theory Exp.* 2005, no. 1, 005, 33 pp. (electronic).
- [12] P. Deift, T. Kriecherbauer, K.T-R. McLaughlin, S. Venakides, and X. Zhou, Uniform Asymptotics for Polynomials Orthogonal with Respect to Varying Exponential Weights and Applications to Universality Questions in Random Matrix Theory, *Commun. Pure Appl. Math.* **52** (1999), 1335–1425.
- [13] P.D. Dragnev and E.B. Saff, Constrained energy problems with applications to orthogonal polynomials of a discrete variable, *J. d'Analyse Math.* **72** (1997), 223–259.
- [14] P.L. Ferrari and H. Spohn, Domino tilings and the six-vertex model at its free fermion point, *J. Phys. A: Math. Gen.* **39** (2006) 1029710306.
- [15] A.G. Izergin, Partition function of the six-vertex model in a finite volume. (Russian) *Dokl. Akad. Nauk SSSR* **297** (1987), no. 2, 331–333; translation in *Soviet Phys. Dokl.* **32** (1987), 878–880.
- [16] A.G. Izergin, D.A. Coker, and V.E. Korepin, Determinant formula for the six-vertex model. *J. Phys. A*, **25** (1992), 4315–4334.
- [17] V. E. Korepin, Calculation of norms of Bethe wave functions. *Commun. Math. Phys.* **86** (1982), 391–418.
- [18] V. Korepin and P. Zinn-Justin, Thermodynamic limit of the six-vertex model with domain wall boundary conditions, *J. Phys. A* **33** No. 40 (2000), 7053
- [19] A.B.J. Kuijlaars, On the finite-gap ansatz in the continuum limit of the Toda lattice, *Duke Math. J.*, **104** (2000) 433–462.
- [20] G. Kuperberg, Another proof of the alternating sign matrix conjecture. *Int. Math. Res. Not.* (1996), 139–150.
- [21] E.H. Lieb, Exact solution of the problem of the entropy of two-dimensional ice. *Phys. Rev. Lett.* **18** (1967) 692.
- [22] E.H. Lieb, Exact solution of the two-dimensional Slater KDP model of an antiferroelectric. *Phys. Rev. Lett.* **18** (1967) 1046–1048.
- [23] E.H. Lieb, Exact solution of the two-dimensional Slater KDP model of a ferroelectric. *Phys. Rev. Lett.* **19** (1967) 108–110.
- [24] E.H. Lieb, Residual entropy of square ice. *Phys. Rev.* **162** (1967) 162.
- [25] E.H. Lieb and F.Y. Wu, Two dimensional ferroelectric models, in *Phase Transitions and Critical Phenomena*, C. Domb and M. Green eds., vol. 1, Academic Press (1972) 331–490.
- [26] E.B. Saff and V. Totik, *Logarithmic potentials with external fields*, Springer, New York, 1997.
- [27] K. Sogo, Toda molecule equation and quotient-difference method. *Journ. Phys. Soc. Japan* **62** (1993), 1887.
- [28] B. Sutherland, Exact solution of a two-dimensional model for hydrogen-bonded crystals. *Phys. Rev. Lett.* **19** (1967) 103–104.
- [29] F.Y. Wu and K.Y. Lin, Staggered ice-rule vertex model. The Pfaffian solution. *Phys. Rev. B* **12** (1975), 419–428.
- [30] E.T. Whittaker and G.N. Watson, *A course of modern analysis*. Fourth edition, reprinted. Cambridge University Press, 2000.
- [31] P. Zinn-Justin, Six-vertex model with domain wall boundary conditions and one-matrix model. *Phys. Rev. E* **62** (2000), 3411–3418.

DEPARTMENT OF MATHEMATICAL SCIENCES, INDIANA UNIVERSITY-PURDUE UNIVERSITY INDIANAPOLIS, 402 N. BLACKFORD ST., INDIANAPOLIS, IN 46202, U.S.A.

E-mail address: bleher@math.iupui.edu

DEPARTMENT OF MATHEMATICAL SCIENCES, INDIANA UNIVERSITY-PURDUE UNIVERSITY INDIANAPOLIS, 402 N. BLACKFORD ST., INDIANAPOLIS, IN 46202, U.S.A.

E-mail address: kliechty@math.iupui.edu